



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>5</sup> :</b> <b>C07D 498/22, 519/00</b> <b>A61K 31/40, 31/55</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/02488</b> <b>(43) International Publication Date:</b> 3 February 1994 (03.02.94)
<b>(21) International Application Number:</b> PCT/US93/06974 <b>(22) International Filing Date:</b> 26 July 1993 (26.07.93)  <b>(30) Priority data:</b> 07/920,102                      24 July 1992 (24.07.92)                      US  <b>(71) Applicants:</b> CEPHALON, INC. [US/US]; 145 Brandywine Parkway, West Chester, PA 19380-4245 (US). KYOWA HAKKO KOGYO CO., LTD. [JP/JP]; Ohtemachi Building, 1-6-1 Ohtemachi, Chiyoda-ku, Tokyo 100 (JP).  <b>(72) Inventors:</b> LEWIS, Michael, E. ; 23 Bills Way, Landenberg, PA 19350 (US). NEFF, Nicola ; 59 Todmorden Drive, Rose Valley, PA 19086 (US). ROBERTS-LEWIS, Jill ; 23 Bills Way, Landenberg, PA 19350 (US). MURAKATA, Chikara ; 2-11-2-702, Bessho, Hachioji-shi, Tokyo 192-03 (JP). SAITO, Hiromitsu ; 5-15-23, Tokura, Mishima-shi, Shizuoka-ken 411 (JP). MATSUDA, Yuzuru ; 1-22-7, Nukuiminami-cho, Koganei-shi, Tokyo 184 (JP). KAUER, James, C. ; 605 Willow Glen Road, Kennett Square, PA 19348 (US).		<b>(74) Agent:</b> CLARK, Paul, T.; Fish & Richardson, 225 Franklin Street, Boston, MA 02110 (US).  <b>(81) Designated States:</b> AU, BR, CA, FI, HU, JP, KR, NO, NZ, PT, RU, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> BIS-STAUROSPORINE AND K-252a DERIVATIVES  <b>(57) Abstract</b>  The invention features novel derivatives of K-252a, as well as novel bis-N-substituted derivatives of staurosporine. Also, a method for treating diseased neuronal cells involving the administration of either the novel staurosporine derivatives or specified functional derivatives of K-252a.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

- 1 -

## "Bis-Staurosporine and K-252a Derivatives"

### Background of the Invention

5 Protein kinases are a broad class of enzymes which act to modify chemically many cellular proteins, by phosphorylation of amino acids.

Inhibitors of protein kinases are structurally varied, and have variable (and sometimes contradictory) effects on the nervous system and other tissues. A given protein kinase inhibitor may influence more than one protein kinase. For example, K-252a, an alkaloid-like material isolated from the culture broth of Nocardiosis sp. and Actinomadula sp. was originally reported to be a protein kinase C inhibitor, but was subsequently found also to inhibit protein kinases A and G, myosin light-chain kinase, and trk (a tyrosine kinase activated by nerve growth factor [NGF], the latter a neurotrophic protein which promotes the survival of peripheral, sensory and sympathetic neurons). Consistent with this latter effect, K-252a blocks the neurotrophic actions of NGF on PC-12 cells (chromaffin cells from rat adrenal medullary tumors, pheochromocytomas), and promotes the survival of dorsal root ganglion neurons and hippocampal neurons. However, it has been found to be cytotoxic at a wide range of concentrations, leading some investigators to conclude that it has limited usefulness in vivo.

A microbial alkaloid related to K-252a, staurosporine, also has a variety of effects on different protein kinases and cell types. Staurosporine was found to have NGF-like effects on PC-12 cells, and to protect the gerbil hippocampus from post-ischemic injury. It is able to reverse damage to cholinergic neurons in the rat basal forebrain.

35 K-252a and staurosporine have been proposed as tumor inhibitors. Staurosporine has been offered as an


- 2 -

insecticide. Derivatives of staurosporine, with a hydrocarbyl radical or an acyl radical substituted at the methylamine nitrogen, have been made and proposed for the following uses: tumor inhibition, inflammation


- 5 inhibition, immunomodulation, and treatment of diseases of the cardiovascular and central nervous systems.

#### Summary of the Invention


The invention features, in one aspect, novel bis-N-substituted derivatives of staurosporine, represented  
10 by the formula



where [Stau] represents a residue of the formula




and W represents a bis(carbamyl) or bis(thiocarbamyl) radical,



where W' is a hydrocarbylene radical of 2-20 carbon atoms and Y is O or S.

The invention also features a novel derivative of K-252a, represented by the formula (II-4):


- 3 -



II-4

where  $R^1$ ,  $R^2$ ,  $Z^1$  and  $Z^2$  are each independently H, X is hydroxymethyl ( $\text{CH}_2\text{OH}$ ), and R is  $\text{OCH}_3$ .

The invention also features a novel derivative of 5 K-252a, represented by the formula:




II-14

where  $R^1$ ,  $R^2$ ,  $Z^1$  and  $Z^2$  are each independently H, X is  $\text{CH}_2\text{-NH-Ser}$ , and R is OH.

Also included in the invention are compounds 10 represented by the following Formula (II-49):


- 4 -



II-49

wherein  $R^2$ ,  $Z^1$ , and  $Z^2$  are each H, R is OH,  $R^1$  is  $\text{CH}_2\text{SO}_2\text{C}_2\text{H}_5$ , and X is  $\text{CO}_2\text{CH}_3$ .


Also included in the invention are compounds  
5 represented by the following Formula (II-38):



II-38

wherein  $R^1$ ,  $R^2$ ,  $Z^1$ , and  $Z^2$  are each H, R is OH, and X is  $\text{CH}_2\text{NHCO}_2\text{C}_6\text{H}_5$

Also included in the invention are compounds  
10 represented by the following Formula (II-45):




II-45

wherein  $R^1$  and  $R^2$  are each Br, R is OH,  $Z^1$  and  $Z^2$  are each H, and X is  $\text{CONHC}_6\text{H}_5$ .

- 5 -


Also included in the invention are compounds represented by the following Formula (II-57):



II-57

wherein  $R^1$ ,  $R^2$ ,  $Z^1$ , and  $Z^2$  are each H, R is OH, and X is  
 5  $\text{CH}_2\text{NHCO}_2\text{CH}_3$ .

Also included in the invention are compounds represented by the following Formula (V):



(V)

in which X represents  $\text{CO}_2\text{R}^5$  (in which  $\text{R}^5$  represents lower  
 10 alkyl) or  $\text{CH}_2\text{NHCO}_2\text{R}^6$  (in which  $\text{R}^6$  represents lower alkyl  
 or aryl);  $\text{R}^1$  represents hydrogen or  $\text{CH}_2\text{SO}_2\text{R}^7$  (in which  $\text{R}^7$   
 represents lower alkyl), provided that the combination of  
 X =  $\text{CO}_2\text{R}^5$  and  $\text{R}^1$  = hydrogen is excluded.

In the definitions of the groups in Formula (V),  
 15 lower alkyl means a straight-chain or branched alkyl  
 group having 1 to 6 carbon atoms, preferably 1 to 3  
 carbon atoms, such as methyl, ethyl, propyl, isopropyl,  
 butyl, isobutyl, sec-butyl, tert-butyl, pentyl,

- 6 -

neopentyl, and hexyl. Aryl means an aryl group having 6 to 10 carbon atoms, such as phenyl and naphthyl.

Formula V compounds can be in the form of pharmaceutically acceptable salts. The pharmaceutically acceptable salts of Compounds (V) include pharmaceutically acceptable acid addition salts, metal salts, ammonium salts, organic amine addition salts, and amino acid addition salts.

Examples of the pharmaceutically acceptable acid addition salts are inorganic acid addition salts such as hydrochloride, sulfate, and phosphate, and organic acid addition salts such as acetate, maleate, fumarate, tartrate, and citrate. Examples of the pharmaceutically acceptable metal salts are alkali metal salts such as sodium salt and potassium salt, alkaline earth metal salts such as magnesium salt and calcium salt, aluminium salt, and zinc salt. Examples of the pharmaceutically acceptable ammonium salts are ammonium salt and tetraethyl ammonium salt. Examples of the pharmaceutically acceptable organic amine addition salts are salts with morpholine and piperidine. Examples of the pharmaceutically acceptable amino acid addition salts are salts with lysine, glycine, and phenylalanine.

In another aspect, the invention features a method for enhancing the function of cholinergic neurons, striatal neurons, and sensory neurons, e.g., dorsal root ganglion neurons, by administering to a mammal, e.g., a human, a therapeutic amount of one of the novel bis-substituted derivatives of staurosporine. The therapy may be given in conjunction with a neurotrophic factor, preferably a member of the neurotrophic family, and most preferably nerve growth factor (NGF). The neurotrophic family is a group of proteins with significant homology to NGF and include, in addition to NGF, brain-derived neurotrophic factor (BDNF; Leibrock et al., Nature






- 7 -

341:149-152, 1989); neutrophil-3 (NT-3; Hohn et al., Nature 344:339-341, 1990); and neurotrophic-5 (NT-5; Berkemeier et al., Neuron 7:857-866, 1991).

In another aspect, the invention features a method for protecting nerve cells of a mammal, e.g., a human, from degeneration induced by excitatory amino acids, by administering to the mammal a therapeutic amount of one of the novel bis-substituted derivatives of staurosporine. Conditions in which such degeneration may occur include Alzheimer's disease; motor neuron disease, e.g., amyotrophic lateral sclerosis; Parkinson's disease; cerebrovascular disease, e.g., ischemic conditions; AIDS dementia; epilepsy; Huntington's disease; and concussive or penetrating injuries to the brain or spinal cord. The therapy may be given in conjunction with a neurotrophic factor, preferably a member of the neurotrophin family, most preferably nerve growth factor (NGF).

In another aspect, the invention features a method for enhancing the function of cholinergic neurons, striatal neurons, and sensory neurons, e.g., dorsal root ganglion neurons, in a mammal, e.g., a human, by administering to the mammal a therapeutic amount of a functional derivative of K-252a, represented by the formulas



25

II

III

or

IV

with any of the following substitutions:

- 8 -

Table 1

	Compound	R <sup>1</sup>	R <sup>2</sup>	X	R	Z <sup>1(1)</sup> Z <sup>2</sup>
5	II-1	H	H	CH <sub>2</sub> N <sub>3</sub>	OH	H
	II-2	NHCONHC <sub>6</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-3	CH <sub>2</sub> SOC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-4	H	H	CH <sub>2</sub> OH	OCH <sub>3</sub>	H
10	II-5	H	H	CONHC <sub>2</sub> H <sub>5</sub>	OH	H
	II-6	H	H	CH=NNH-	OH	H
	II-7 (2,7)	H	H	CH <sub>2</sub> NH-Gly	OH	H
	II-8	H	H	CON(CH <sub>3</sub> ) <sub>2</sub>	OH	H
	II-9 (3)	H	H	-CH <sub>2</sub> NHCO <sub>2</sub> -		H
15	II-10	Br	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-11	H	H	CONH <sub>2</sub>	OH	H
	II-12	H	H	CH <sub>2</sub> OH	OH	H
	III-1	--	H	--	--	H
	II-13	H	H	CONHC <sub>3</sub> H <sub>7</sub>	OH	H
20	II-14 (2)	H	H	CH <sub>2</sub> NH-Ser	OH	H
	II-15	H	H	CH <sub>2</sub> SOCH <sub>3</sub>	OH	H
	II-16	H	H	CH=NOH	OH	H
	II-17	H	H	CON O	OH	H
	II-18 (2,7)	H	H	CH <sub>2</sub> NH-Pro	OH	H
25	II-19	H	H	CH=NNHC(=NH)NH <sub>2</sub>	OH	H
	II-20	Br	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	O
	II-21	H	H	CONH(CH <sub>2</sub> ) <sub>2</sub> OH	OH	H
	II-22	H	H	CO <sub>2</sub> CH <sub>3</sub>	OH	O
	III-2	--	H	--	--	O
30	II-23	H	H	H	OH	H
	II-24	H	H	CH=NNHCONH <sub>2</sub>	OH	H
	II-25	H	H	CH <sub>2</sub> OCOCH <sub>3</sub>	OH	H
	II-26 (3)	H	H	-CH <sub>2</sub> OC(CH <sub>3</sub> ) <sub>2</sub> O-		H
	II-29	NHCONHC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
35	II-30	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-31	Br	H	CH <sub>2</sub> OH	OH	H
	II-32	Br	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-33	CH <sub>2</sub> SC <sub>6</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-34	Cl	Cl	CO <sub>2</sub> CH <sub>3</sub>	OH	H
40	II-36	H	H	CONHC <sub>6</sub> H <sub>5</sub>	OH	H
	II-37	H	H	CH <sub>2</sub> SO	OH	H
	II-38	H	H	CH <sub>2</sub> NHCO <sub>2</sub> C <sub>6</sub> H <sub>5</sub>	OH	H
	II-39	NHCONHC <sub>2</sub> H <sub>5</sub>	NHCONHC <sub>2</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-40	N(CH <sub>3</sub> ) <sub>2</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
45	II-41	CH <sub>3</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-42	CH <sub>2</sub> OCONHC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-43	NHCO <sub>2</sub> CH <sub>3</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-44	Br	Br	CH <sub>2</sub> OH	OH	H
	II-45	Br	Br	CONHC <sub>6</sub> H <sub>5</sub>	OH	H
50	II-46	Br	Br	CONHCH <sub>2</sub> CH <sub>2</sub> OH	OH	H
	II-47	CH <sub>2</sub> OC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-48	CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H

- 9 -


	II-49	CH <sub>2</sub> SO <sub>2</sub> C <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-50	CH <sub>2</sub> S	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-51	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-52	CH=NNH	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
5	II-53	CH <sub>2</sub> S	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-54	CH <sub>2</sub> S(O)	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-55	CH <sub>2</sub> S(O)	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-56	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> OH	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-57	H	H	CH <sub>2</sub> NHCO <sub>2</sub> CH <sub>3</sub>	OH	H
10	II-58	Br	H	CONH <sub>2</sub>	OH	H
	II-59	H	H	CH <sub>2</sub> SC <sub>6</sub> H <sub>5</sub>	OH	H
	II-60	H	H	CH <sub>2</sub> S	OH	H
	II-61	H	H	CH <sub>2</sub> SOC <sub>6</sub> H <sub>5</sub>	OH	H
	IV-1 <sup>(4,9)</sup>	H	H	--	--	H
15	IV-2 <sup>(5)</sup>	Br	H	--	--	H
	IV-3 <sup>(6)</sup>	H	H	--	--	H
	IV-4 <sup>(8,9)</sup>	H	H	--	--	H

- (1) Z<sup>1</sup> and Z<sup>2</sup> are both hydrogen, or both are  
 20 combined together to represent oxygen, where indicated.  
 (2) NH-amino acid linkage is an amide bond  
 through the carboxyl group of the amino acid.  
 (3) X and R are combined together to form the  
 linking group.  
 25 (4) R<sup>3</sup> is CH<sub>2</sub>CH=CH<sub>2</sub>; R<sup>4</sup> is H.  
 (5) R<sup>3</sup> and R<sup>4</sup> are each H.  
 (6) R<sup>3</sup> and R<sup>4</sup> are each CH<sub>2</sub>CH=CH<sub>2</sub>.  
 (7) Compound is in the form of the  
 hydrochloride.  
 30 (8) R<sup>3</sup> is H and R<sup>4</sup> is CH<sub>2</sub>CH=CH<sub>2</sub>.  
 (9) IV-1 and IV-4 is a 1.5 to 1.0 mixture of  
 the two components.


The therapy may be given in conjunction with a  
 35 neurotrophic factor, preferably a member of the  
 neurotrophin family, most preferably nerve growth factor  
 (NGF). The method may be used to treat Huntington's  
 disease.

In a preferred aspect, the invention features  
 40 a method for enhancing the function of a dorsal root  
 ganglion nerve cell, by administering to a mammal, e.g.,  
 a human, a therapeutic amount of a functional derivative  
 of K-252a, represented by the formula (II) or (III)

- 10 -



(II)



(III)

wherein the following substitutions are made:

Table 2

5	Compound <sup>(1)</sup>	R <sup>1</sup>	X	R	Z <sup>1</sup> (2) Z <sup>2</sup>
	II-1	H	CH <sub>2</sub> N <sub>3</sub>	OH	H
	II-2	NHCONHC <sub>6</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
10	II-3	CH <sub>2</sub> SOC <sub>2</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-4	H	CH <sub>2</sub> OH	OCH <sub>3</sub>	H
	II-5	H	CONHC <sub>2</sub> H <sub>5</sub>	OH	H
	II-6	H	CH=NNH-	OH	H
	II-8	H	CON(CH <sub>3</sub> ) <sub>2</sub>	OH	H
15	II-9 <sup>(3)</sup>	H	-CH <sub>2</sub> NHCO <sub>2</sub> -		H
	II-10	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-11	H	CONH <sub>2</sub>	OH	H
	II-12	H	CH <sub>2</sub> OH	OH	H
	III-1	--	--	--	H
20	II-13	H	CONHC <sub>3</sub> H <sub>7</sub>	OH	H
	II-15	H	CH <sub>2</sub> SOCH <sub>3</sub>	OH	H
	II-17	H	CON <sup>O</sup>	OH	H
	II-19	H	CH=NNHC(=NH)NH <sub>2</sub>	OH	H
	II-20 <sup>(1)</sup>	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	O
25	II-21	H	CONH(CH <sub>2</sub> ) <sub>2</sub> OH	OH	H
	III-2	--	--	--	O
	II-23	H	H	OH	H
	II-24	H	CH=NNHCONH <sub>2</sub>	OH	H
	II-25	H	CH <sub>2</sub> OCOCH <sub>3</sub>	OH	H
30	II-30	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-32	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	H

(1) R<sup>2</sup> is hydrogen, except in compound II-20 and II-32 where R<sup>2</sup>=Br.


35 (2) Z<sup>1</sup> and Z<sup>2</sup> are both hydrogen, or both are combined together to represent oxygen, where indicated.

- 11 -

(3) X and R are combined together to form the linking group.

The therapy may be given in conjunction with a neurotrophic factor, preferably a member of the neurotrophin family, most preferably nerve growth factor (NGF).

In a preferred aspect, the invention features a method for enhancing the function of cholinergic neurons of a mammal, e.g., a human, by administering to the mammal a therapeutic amount of K-252a, represented by the formula (II):




(II)


wherein  $R^1$  and  $R^2$  are each H, X is  $\text{CO}_2\text{CH}_3$ , R is OH, and  $Z^1$  and  $Z^2$  are each H. The therapy may be given in conjunction with a neurotrophic factor, preferably a member of the neurotrophin family, most preferably nerve growth factor (NGF), and may be used to treat Huntington's disease.

In a preferred aspect, the invention features a method for enhancing the survival and/or function of a striatal nerve cell, by administering to a mammal, e.g., a human, a therapeutic amount of K-252a or a functional derivative of K-252a, represented by the formula (II), (III), or (IV):


- 12 -



II



III



IV

wherein the following substitutions are made:

Table 3

5	Compound	R <sup>1</sup>	R <sup>2</sup>	X	R	$\begin{matrix} Z^2(1) \\ Z^1 \end{matrix}$
	K-252a	H	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	III-1	--	H	--	--	H
	II-1	H	H	CH <sub>2</sub> N <sub>3</sub>	OH	H
10	II-35	H	H	CO <sub>2</sub> n-C <sub>6</sub> H <sub>13</sub>	OH	H
	II-20	Br	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	O
	II-10	Br	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-28	O-n-C <sub>3</sub> H <sub>7</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-5	H	H	CONHC <sub>2</sub> H <sub>5</sub>	OH	H
15	II-29	NHCONHC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-2	NHCONHC <sub>6</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-3	CH <sub>2</sub> SOC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-30	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-6	H	H	CH=NNH	OH	H
20	II-31	Br	H	CH <sub>2</sub> OH	OH	H
	II-32	Br	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	IV-1 <sup>(2)</sup>	--	H	--	--	H
	II-33	CH <sub>2</sub> SC <sub>6</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-34	Cl	Cl	CO <sub>2</sub> CH <sub>3</sub>	OH	H
25						

(1) Z<sup>1</sup> and Z<sup>2</sup> are both hydrogen, or both are combined together to represent oxygen, where indicated.

(2) R<sup>3</sup> is CH<sub>2</sub>-CH=CH<sub>2</sub>; R<sup>4</sup> is H.

The therapy may be given in conjunction with a  
 30 neurotrophic factor, preferably a member of the  
 neurotrophin family, most preferably nerve growth factor,  
 and may be used to treat Huntington's disease.

- 13 -

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

Description of the Preferred Embodiments

5           The drawings are first described.

Drawings

Fig. 1 is a graph illustrating the effect of K-252a, 1,6-hexamethylene-bis-(carbamylstaurosporine) (HBCS), and staurosporine on basal ornithine  
10 decarboxylase (ODC) activity in PC-12 cells.

Fig. 2 is a graph illustrating the effects of staurosporine, HBCS, and K-252a on NGF-stimulated ODC activity in PC-12 cells.

Fig. 3 is a graph illustrating the NGF-  
15 potentiating effect of HBCS on ODC activity in PC-12 cells.

Fig. 4 is a graph illustrating the effect of K-252a on choline acetyltransferase (ChAT) specific activity in rat embryonic spinal cord cultures.

20           Fig. 5 is a graph illustrating the time course of K-252a effect on ChAT activity in rat embryonic spinal cord cultures.

Fig. 6 is a graph illustrating the effect of K-252a on survival of chick embryonic dorsal root ganglion  
25 neurons.

Fig. 7 is a graph illustrating the effect of K-252a functional derivatives on survival of chick embryonic dorsal root ganglion neurons.

Fig. 8 is a graph illustrating the effect of K-  
30 252a functional derivatives on ChAT activity in rat embryonic spinal cord cultures.

Fig. 9 is a graph illustrating the effect of K-252a on kainate-induced damage to the rat hippocampus.

- 14 -

Fig. 10 is a graph illustrating the effect of K-252a on kainate-induced spectrin proteolysis in the rat hippocampus.

Fig. 11 is a graph illustrating the effect of HBCS  
5 on kainate-induced damage to the hippocampus.

Fig. 12 is a graph illustrating the effect of K-252a analogs on kainate-induced spectrin proteolysis in the rat hippocampus.

Fig. 13 is a table showing the relative activity  
10 of K-252a derivatives on ChAT activity in rat spinal cord cultures.

Fig. 14 is a table showing the relative activity of K-252a derivatives on neuronal survival in chick dorsal root ganglion cultures.

Fig. 15 is a graph illustrating survival of  
15 striatal neurons in the presence of K-252a.

Fig. 16 is a graph illustrating survival of striatal cells in the presence of K-252a.

Fig. 17 is a photomicrograph of striatal neurons  
20 cultured in the presence or absence of K-252a.

Fig. 18 is a table showing the relative activity of K-252a derivatives on neuronal survival in rat striatal cultures.

Fig. 19 shows the synthesis of Compound H from  
25 starting Compound C.

Fig. 20 shows the synthesis of Compound II-45 from starting Compound J.

#### Staurosporine Derivatives


The present invention relates to novel bis-N-  
30 substituted derivatives of staurosporine and their use as therapeutics for neurological diseases, especially those diseases characterized either by neuronal cells which are injured, compromised, undergoing axonal degeneration, or at increased risk of dying, or by impaired cholinergic  
35 activity. These diseases include those induced by




- 15 -

excitatory amino acids. The therapeutic use of these novel derivatives includes use of the derivatives alone and use of the derivatives combined with exogenous administration of neurotrophic factors (preferably

5 members of the neurotrophin family, most preferably nerve growth factor, NGF). The compounds within the scope of this invention may be represented by the formula



in which [Stau] represents a residue of the formula:



10 and W represents a bis(carbamyl) or bis(thiocarbamyl) radical,



in which W' is a hydrocarbylene radical of 2-20 carbon atoms and Y is O or S. W' is preferably an alkylene

15 radical of 2-10 carbons, unsubstituted, or substituted with 1-3 alkyl groups of 1-3 carbons; an arylene radical of 6-12 carbons, unsubstituted, or substituted with 1-3 alkyl groups of 1-3 carbons, chlorine or bromine. W' is especially preferably hexamethylene and 1,4-phenylene. Y

20 is preferably O.

Compounds of formula (I) can be prepared by procedures known in the art for preparation of carbamates and thiocarbamates. Preferably, the compounds are prepared by reaction of a bis-diisocyanate or a bis-

25 diisothiocyanate with staurosporine to give a compound of formula (I) wherein Y=O or Y=S respectively.

- 16 -

Intermediate bis-diisocyanates and bis-diisothiocyanates suitable for use include:

- 1,6-diisocyanatohexane
- toluene-2,6-diisocyanate
- 5 benzene-1,2-diisocyanate
- 2-methyl-1,5-diisocyanatopentane
- naphthalene-2,6 diisocyanate
- 1,6-diisothiocyanatohexane
- 1,4-diisothiocyanatobutane
- 10 toluene-2,4-diisocyanate
- benzene-1,4-diisocyanate
- 1,2-diisocyanatoethane
- naphthalene-1,5-diisocyanate
- 1,5-diisocyanatopentane
- 15 benzene-1,4-diisothiocyanate
- 2-methyl-1,5-diisothiocyanatopentane

For reviews of the preparation of isocyanates and isothiocyanates, see the article by Richter and Ulrich, pp. 619-818, in Patai, "The Chemistry of Cyanates and  
20 Their Thio Derivatives", Part 2, Wiley, New York, 1977. The compounds are preferably prepared by reaction of phosgene (Y=O) or thiophosgene (Y=S) with the corresponding diamine. Alternative methods of preparation may also be employed. For example, 1,2-  
25 diisocyanatoethane may be prepared by reaction of ethylene urea with phosgene followed by heating.


#### K-252a Derivatives

The present invention is also directed to the use of specific functional derivatives of K-252a, as  
30 therapeutics in certain neurological diseases or disturbances characterized by neurons which are injured, compromised, undergoing axonal degeneration, or at risk of dying. The functional derivatives may be administered alone or in conjunction with a neurotrophic factor  
35 (preferably a member of the neurotrophin family, most preferably nerve growth factor, NGF). A "functional derivative" of K-252a is defined as a modified form of that molecule, which possesses the desired biological activity, herein defined as neuroprotective activity, for


- 17 -

example the ability to promote nerve cell survival, or to promote nerve fiber (e.g. axonal) growth, or to enhance cholinergic nerve cell function, or to enhance the function of sensory cells, e.g., dorsal root ganglion nerve cells, or to enhance the function and/or survival of striatal neurons. Such molecular modifications may improve the molecule's solubility, absorption, transport (e.g., through the blood-brain barrier and cellular membranes), biological half-life, etc. Alternatively, or in addition, some moieties may decrease the toxicity of the molecule, or eliminate or attenuate any undesirable side effect of the molecule.


The compounds within the scope of the invention may be represented by formula (II) [hereinafter referred to as compound (II)], formula (III) [hereinafter referred to as compound (III)], and formula (IV) [hereinafter referred to as compound (IV)], below:



(II)



(III)



(IV)

with substitutions in Table 4, below, depicting the compounds within the scope of the invention. The functional derivatives of K-252a of the invention may be prepared *de novo* by chemical synthesis using methods known to those skilled in the art. For example, procedures used for preparation of Compound II are described by Murakata et al (U.S. Patent 4,923,986),

- 18 -

hereby incorporated by reference. Procedures used for preparation of Compound III are described by Moody et al., J. Org. Chem. 57: 2105-2114 (1992); Steglich et al., Angew. Chem. Int. Ed. Engl. 19: 459-460 (1980); Nakanishi et al., J. Antibiotics 39: 1066-1071 (1986); and Japanese Patent Application No. 60-295172 (1985). Further methods are described for compounds II-1, 9, 12 and 15 in Japanese Patent Application No. 60-295173 (1985); compounds II-2, 3, 4, 24, 25 and 26 in Japanese Patent Application No. 62-327858 (1987); compounds II-20 in Japanese Patent Application No. 62-327859 (1987); and compounds II-10 in Japanese Patent Application No. 60-257652 (1985) by Meiji Seika Kaisha Ltd.

Table 4: Functional Derivatives of K-252a<sup>(10)</sup>

Compound	R <sup>1</sup>	R <sup>2</sup>	X	R	Z <sup>1(1)</sup> Z <sup>2</sup>
II-1	H	H	CH <sub>2</sub> N <sub>3</sub>	OH	H
II-2	NHCONHC <sub>6</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
20 II-3	CH <sub>2</sub> SOC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
II-4	H	H	CH <sub>2</sub> OH	OCH <sub>3</sub>	H
II-5	H	H	CONHC <sub>2</sub> H <sub>5</sub>	OH	H
II-6	H	H	CH=NNH-	OH	H
II-7(2,7)	H	H	CH <sub>2</sub> NH-Gly	OH	H
25 II-8	H	H	CON(CH <sub>3</sub> ) <sub>2</sub>	OH	H
II-9(3)	H	H	-CH <sub>2</sub> NHCO <sub>2</sub> -		H
II-10	Br	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
II-11	H	H	CONH <sub>2</sub>	OH	H
II-12	H	H	CH <sub>2</sub> OH	OH	H
30 III-1	--	H	--	--	H
II-13	H	H	CONHC <sub>3</sub> H <sub>7</sub>	OH	H
II-14(2)	H	H	CH <sub>2</sub> NH-Ser	OH	H
II-15	H	H	CH <sub>2</sub> SOCH <sub>3</sub>	OH	H
II-16	H	H	CH=NOH	OH	H
35 II-17	H	H	CON O	OH	H
II-18(2,7)	H	H	CH <sub>2</sub> NH-Pro	OH	H
II-19	H	H	CH=NNHC(=NH)NH <sub>2</sub>	OH	H
II-20	Br	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	O
II-21	H	H	CONH(CH <sub>2</sub> ) <sub>2</sub> OH	OH	H
40 II-22	H	H	CO <sub>2</sub> CH <sub>3</sub>	OH	O
III-2	--	H	--	--	O
II-23	H	H	H	OH	H
II-24	H	H	CH=NNHCONH <sub>2</sub>	OH	H
II-25	H	H	CH <sub>2</sub> OCOCH <sub>3</sub>	OH	H
45 II-26(3)	H	H	-CH <sub>2</sub> OC(CH <sub>3</sub> ) <sub>2</sub> O-		H

- 19 -

	II-29	NHCONHC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-30	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-31	Br	H	CH <sub>2</sub> OH	OH	H
	II-32	Br	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	H
5	II-33	CH <sub>2</sub> SC <sub>6</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-34	Cl	Cl	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-36	H	H	CONHC <sub>6</sub> H <sub>5</sub>	OH	H
	II-37	H	H	CH <sub>2</sub> SO	OH	H
	II-38	H	H	CH <sub>2</sub> NHCO <sub>2</sub> C <sub>6</sub> H <sub>5</sub>	OH	H
10	II-39	NHCONHC <sub>2</sub> H <sub>5</sub>	NHCONHC <sub>2</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-40	N(CH <sub>3</sub> ) <sub>2</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-41	CH <sub>3</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-42	CH <sub>2</sub> OCONHC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-43	NHCO <sub>2</sub> CH <sub>3</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
15	II-44	Br	Br	CH <sub>2</sub> OH	OH	H
	II-45	Br	Br	CONHC <sub>6</sub> H <sub>5</sub>	OH	H
	II-46	Br	Br	CONHCH <sub>2</sub> CH <sub>2</sub> OH	OH	H
	II-47	CH <sub>2</sub> OC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-48	CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
20	II-49	CH <sub>2</sub> SO <sub>2</sub> C <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-50	CH <sub>2</sub> S	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-51	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-52	CH=NNH	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-53	CH <sub>2</sub> S	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
25	II-54	CH <sub>2</sub> S(O)	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-55	CH <sub>2</sub> S(O)	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-56	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> OH	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-57	H	H	CH <sub>2</sub> NHCO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-58	Br	H	CONH <sub>2</sub>	OH	H
30	II-59	H	H	CH <sub>2</sub> SC <sub>6</sub> H <sub>5</sub>	OH	H
	II-60	H	H	CH <sub>2</sub> S	OH	H
	II-61	H	H	CH <sub>2</sub> SOC <sub>6</sub> H <sub>5</sub>	OH	H
	IV-1(4,9)	H	H	--	--	H
	IV-2(5)	Br	H	--	--	H
35	IV-3(6)	H	H	--	--	H
	IV-4(8,9)	H	H	--	--	H

(1) Z<sup>1</sup> and Z<sup>2</sup> are both hydrogen, or both are combined together to represent oxygen, where indicated.

40 (2) NH-amino acid linkage is an amide bond through the carboxyl group of the amino acid.

(3) X and R are combined together to form the linking group.

(4) R<sup>3</sup> is CH<sub>2</sub>CH=CH<sub>2</sub>; R<sup>4</sup> is H.

45 (5) R<sup>3</sup> and R<sup>4</sup> are each H.

(6) R<sup>3</sup> and R<sup>4</sup> are each CH<sub>2</sub>CH=CH<sub>2</sub>.

(7) Compound is in the form of the hydrochloride.

- 20 -

(8)  $R^3$  is H and  $R^4$  is  $\text{CH}_2\text{CH}=\text{CH}_2$ .

(9) IV-1 and IV-4 is a 1.5 to 1.0 mixture of the two components.

(10) For K-252a itself,  $R^1=R^2=\text{H}$ ,  $\text{X}=\text{CO}_2\text{CH}_3$ ,  
5  $\text{R}=\text{OH}$ ,  $\text{Z}^1$  and  $\text{Z}^2=\text{H}$ .

The invention also involves a method for enhancing the function of cholinergic neurons, by administration of a therapeutic amount of K-252a, represented by the formula (II) given above and  
10 substitutions shown in Table 4, (note 10). This compound is prepared by procedures described in the art (see Matsuda et al., U.S. Patent 4,554,402; Kase et al., J. Antibiotics 37: 1059-1065 [1986]). By "enhancing the  
15 function of cholinergic neurons" is meant promoting cholinergic nerve cell survival, and/or nerve fiber (e.g. axonal) growth, and/or enhancing cholinergic activity of nerve cells. K-252a may be administered with or without a neurotrophic factor, preferably a member of the  
20 neurotrophin family, most preferably nerve growth factor (NGF).

#### Uses of the Compounds

As described more fully below, the present invention provides novel uses of functional derivatives  
25 of K-252a or compounds of Formula I, either alone or in combination with neurotrophic factors such as NGF, as therapeutics for neurological diseases, especially those diseases characterized either by neuronal cells which are injured, compromised, undergoing axonal degeneration, or  
30 at increased risk of dying, or by impaired cholinergic activity. These diseases include those induced by excitatory amino acids. The bioactivity of the compounds of the invention, including the combination with a neurotrophic factor, may conveniently be assayed by a  
35 cultured PC-12 cell ornithine decarboxylase assay, a cultured spinal cord choline acetyltransferase assay, a cultured dorsal root ganglion neuron survival assay, a

- 21 -

cultured striatal neuron survival assay, or an *in vivo* excitotoxin neuroprotection assay, all of which are described in detail below. Thus, the compounds of this invention are useful for administration to humans or  
5 other mammals who suffer from neurological diseases or disturbances characterized by increased risk of neuronal cell death or dysfunction, as described above. These neurological diseases and disturbances include but are not limited to: Alzheimer's disease; motor neuron disease  
10 including amyotrophic lateral sclerosis; Parkinson's disease; stroke or other ischemic injuries; Huntington's disease; AIDS dementia; epilepsy; concussive or penetrating injuries of the brain or spinal cord; and peripheral neuropathies.

15 The compounds provided herein can be formulated into pharmaceutical compositions by admixture with pharmaceutically acceptable nontoxic excipients and carriers. As noted above, such compositions may be prepared for use in parenteral administration,  
20 particularly in the form of liquid solutions or suspensions; for oral administration, particularly in the form of tablets or capsules; or intranasally, particularly in the form of powders, nasal drops, or aerosols.

25 The composition may conveniently be administered in unit dosage form and may be prepared by any of the methods well known in the pharmaceutical art, for example, as described in Remington's Pharmaceutical Sciences (Mack Pub. Co, Easton, PA, 1980). Formulations  
30 for parenteral administration may contain as common excipients sterile water or saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes and the like. In particular, biocompatible, biodegradable lactide polymer,  
35 lactide/glycolide copolymer, or polyoxyethylene-

- 22 -

polyoxypropylene copolymers may be useful excipients to control the release of the active compounds. Other potentially useful parenteral delivery systems for these active compounds include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation administration contain as excipients, for example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or oily solutions for administration in the form of nasal drops, or as a gel to be applied intranasally. Formulations for parenteral administration may also include glycocholate for buccal administration, methoxysalicylate for rectal administration, or citric acid for vaginal administration.

The materials of this invention can be employed as the sole active agent in a pharmaceutical or can be used in combination with other active ingredients, e.g., other growth factors which could facilitate neuronal survival or axonal growth in neurological diseases or disorders, for example, peripheral neuropathy.

The concentrations of the compounds described herein in a therapeutic composition will vary depending upon a number of factors, including the dosage of the drug to be administered, the chemical characteristics (e.g., hydrophobicity) of the compounds employed, and the route of administration. In general terms, the compounds of this invention may be provided in an aqueous physiological buffer solution containing about 0.1 to 10% w/v compound for parenteral administration. Typical dose ranges are from about 1  $\mu$ g/kg to about 1 g/kg of body weight per day; a preferred dose range is from about 0.01 mg/kg to 100 mg/kg of body weight per day. The preferred dosage of drug to be administered is likely to depend on



- 23 -

such variables as the type and extent of progression of the neurological disease, the overall health status of the particular patient, the relative biological efficacy of the compound selected, the formulation of the compound  
5 excipients, and its route of administration.

The present invention will be further illustrated by the following examples. These examples are not to be construed as limiting the scope of the invention, which is to be determined solely by the  
10 appended claims.

#### Example 1

##### 1,6-Hexamethylene-bis-(carbamylstaurosporine) (HBCS)

A solution of 1.0 mg (2.15 micromoles) of staurosporine (Kamiya Biomedical Company, Thousand Oaks,  
15 CA) in 1.00 ml of ethyl acetate (dried over anhydrous magnesium sulfate) was treated with 17 microliters (1.08 micromoles) of a solution of 10.75 mg of hexamethylene-bis-isocyanate in 1.0 ml of dried ethyl acetate. The reaction mixture in an amber glass reaction vial was  
20 allowed to stand at room temperature for two days. A crystalline deposit weighing 600 micrograms was separated. Its composition was verified by fast atom bombardment mass spectroscopy (FAB-MS).

25       M+H<sup>+</sup> Calculated = 1102       M+Na<sup>+</sup> Calculated = 1124  
                                Found = 1102                                  Found = 1124

This product and all of the subsequently described staurosporine derivatives were stored in non-actinic glass vials.

#### Example 2

##### 30 p-Phenylene-bis-(carbamylstaurosporine) (PBCS)

A solution of 1.0 mg. of staurosporine (2.15 micromoles) in 1.00 ml of dried ethyl acetate was treated with 45 microliters (1.08 micromoles) of a solution prepared from 3.83 mg of p-phenylene diisocyanate (Trans

- 24 -

World Chemicals P1586-1) in 1.00 ml of dried ethyl acetate. The reaction mixture was allowed to stand overnight. A white precipitate deposited. Then 0.5 ml of petroleum ether was added. The mixture was filtered into a vacuum-dried sintered-glass funnel. A total of 0.90 mg of crystalline product was collected and was identified as p-phenylene-bis-(carbamylstaurosporine) by fast atom bombardment mass spectroscopy.

10  $M+H^+$  Calculated = 1093  
Found = 1093

#### Preparation A

##### N-Phenylcarbamylstaurosporine (PCS)

Reference: U.S. Patent 5,093,330

15 A solution of 2.0 mg of staurosporine (4.30 micromoles) in 1.50 ml of dried ethyl acetate was treated with 468  $\mu$ l (4.30 micromoles) of a solution of 10  $\mu$ l of phenyl isocyanate in 0.990 ml of dried ethyl acetate. The solution was allowed to stand overnight and 3 ml of  
20 hexade was added in portions. Colorless crystals were obtained which weighed 2.39 mg. After recrystallizing this product from 1 ml of ethyl acetate and 2 ml of petroleum ether, 1.75 mg of a crystalline product was isolated. From a similar preparation, the product's  
25 composition as N-phenylcarbamylstaurosporine was verified by FAS-MS.

$M+H^+$  Calculated = 586  
Found = 586

#### Preparation B

##### N-Phenylthiocarbamylstaurosporine (PTCS)

30 A solution of 1.0 mg (2.15 micromoles) of staurosporine in 1.00 ml of ethyl acetate was treated with 26 microliters of a stock solution of 10 microliters of phenyl isothiocyanate in 1.00 ml of ethyl acetate.  
35 This aliquot contained 290 micrograms (2.15 micromoles)

- 25 -

of phenyl isothiocyanate. The reaction mixture was held at 25°C overnight, and then 2.0 ml of hexade was added. The resulting crystalline product was filtered off, washed with hexade and dried with a stream of argon gas.

5                   FAB-MS           Calc:  $M+H^+ = 602$   
                                  Found           = 602

### Preparation C

#### N-Ethylcarbamyldaurosporine (ECS)

A solution of 0.9 mg (1.93 micromoles) of  
10 staurosporine in 900 microliters of ethyl acetate was  
treated with 1.93 micromoles (30.2 microliters of a stock  
solution of 9.05 mg of ethyl isocyanate in 2.00 ml of  
dried ethyl acetate) of ethyl isocyanate. The reaction  
mixture was held at 25°C overnight, and 2.0 ml of hexade  
15 was added. The crystalline product was separated and  
dried.

FAB-MS   Calc.:  $M+H^+ = 538$             $M+Na^+ = 560$   
          Found           = 538               = 560

### Example 3

#### 20 Compound II-4


Compound A (962 mg, 2 mmol) was dissolved in a  
mixture of 30 ml of tetrahydrofuran and 10 ml of  
methanol, and then 760 mg of sodium borohydride (20 mmol)  
was added thereto under ice cooling, followed by stirring  
25 at the same temperature for 4 hours and further at room  
temperature for 12 hours. After 3N hydrochloric acid was  
added thereto, the solution was washed with an aqueous  
solution of sodium chloride and dried over magnesium  
sulfate, followed by evaporation of the solvent. The  
30 residue was purified by silica gel column chromatography  
(chloroform/methanol = 98/2) to give 882 mg (yield 97%)  
of Compound II-4.

Melting Point: 130-140°C

$^1H$ -NMR (DMSO- $d_6$ )  $\delta$  (ppm): 2.032 (1H, dd, J=5.0,

- 26 -

13.9Hz), 2.231 (3H, s), 2.967 (3H, s), 3.609 (1H, dd, J=7.6, 13.4Hz), 3.959 (2H, m), 5.000 (2H, s), 5.268 (1H, t, J=5.3Hz), 7.065 (1H, dd, J=4.9, 7.3Hz), 7.254-8.038 (7H, m), 8.565 (1H, s), 9.206 (1H, d, J=7.8Hz)

Compound AExample 410 Compound II-14

Compound B (393 mg, 0.9 mmol) was dissolved in 25 ml of tetrahydrofuran, and then 3 ml of tetrahydrofuran containing 309 mg of carbobenzoxy-L-serine (1.35 mmol), 156 mg of N-oxysuccinimide (1.35 mmol), 0.1 ml of 4-methylmorpholine (0.9 mmol) and 279 mg of dicyclohexylcarbodiimide (1.35 mmol) was added under ice cooling, followed by stirring for 12 hours. The reaction mixture was filtered and the solvent was evaporated. The residue was purified by silica gel column chromatography (chloroform/methanol = 99/1) to give 429 mg (yield 72%) of Compound C.

Melting Point: 188-193°C

SIMS (m/z): 660 (M+1)<sup>+</sup>

Compound C (399 mg) was dissolved in 10 ml of 25 dimethylformamide, and then 300 mg of 10% palladium on

- 27 -

carbon was added, followed by stirring at 50°C for 7 hours in a hydrogen stream. The reaction mixture was filtered through Celite and the solvent was evaporated. The residue was purified by silica gel column


5 chromatography (chloroform/methanol/28% ammonium hydroxide = 90/10/1) and the obtained product was dissolved in 5 ml of tetrahydrofuran, followed by addition of 5 ml of 1.7N hydrogen chloride/ethyl acetate and 10 ml of diethyl ether. The precipitate was  
10 separated from the solution by filtration to give 234 mg (yield 69%) of Compound II-14.

Melting Point: >300°C


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub> + D<sub>2</sub>O) δ (ppm): 1.92-2.28(1H, m),  
2.20 (3H, s), 2.84-3.12(7H, m),  
15 3.40-4.20(5H, m), 5.04 (2H, s),  
6.98(1H, m), 7.24-8.20(7H, m),  
8.76(1H, brs), 9.22(1H, d, J=8Hz)

SIMS (m/z): 527 (M+2)<sup>+</sup>

Compound B



Compound C



CbZ: carbobenzoxy

20

Example 5

PC-12 cells are a clonal population arising from a tumor of rat adrenal medulla, and have proven to be an extremely useful and widely studied model for study of the actions of NGF (Guroff, Cell Culture in the

SUBSTITUTE SHEET

- 28 -

Neurosciences, Plenum Publishing Corporation, 1985, Chapter 8, pages 245-272). One particularly robust effect of NGF on these cells is a rapid stimulation of the activity of ornithine decarboxylase (ODC), an effect  
5 which was reported to be blocked by 200 nM K-252a (Koizumi et al., 1988). In the experiments of this Example, PC-12 cells (obtained from Dr. G. Guroff) were cultured in 48-well plates at a density of  $6 \times 10^4$  cells/cm<sup>2</sup> and incubated with drug vehicle (0.5% DMSO),  
10 staurosporine, or HBCS. K-252a and staurosporine are commercially available from Kamiya Biomedical. Four hours after drug addition, the cells were harvested for ODC assay, as described by Huff et al. (J. Cell Biol. 88: 189-198, 1981).

15 All three compounds produced an induction (i.e. increase) of ODC activity, but there were considerable differences in potency and efficacy (Fig. 1). K-252a produced a dose-dependent induction of ODC activity, with effects detectable at 2 nM and increasing  
20 to a maximum at 200 nM (36.3 fold induction). The effects of staurosporine were likewise detectable at 2 nM, but peaked at 20 nM (34.7 fold induction), and declined considerably at 200 nM. HBCS similarly induced at 2 nM, but higher concentrations failed to yield an  
25 increased effect, so that the maximum efficacy was much less than that of the other two compounds (6.5 fold induction). In another experiment, the effects of PTCS, PCS, and ECS on PC-12 cell ODC activity were compared to that of K-252a. At 200 nM concentrations, expressing the  
30 activity of K-252a as 100%, PTCS exhibited 71.4% of the activity of K-252a, while PCS and ECS exhibited 88.9% and 61.9% of the activity of K-252a, respectively. However, the protein kinase C inhibitor H-7 did not induce ODC activity at 30  $\mu$ M, a concentration known to inhibit

- 29 -

protein kinase C activity (Nakadate et al., Biochem. Pharmacol. 37: 1541-1545, 1988).

The ability of K-252a, staurosporine and HBCS to potentiate and/or inhibit NGF bioactivity was assessed by adding 10 ng NGF per ml of cell culture medium, in the absence or presence of the above compounds in the concentrations previously indicated, followed by ODC assay of the cells as above (Fig. 2). This concentration of NGF was selected to provide an intermediate level of induction so that either potentiating or inhibiting effects of the compounds could be detected. K-252a at 200 nM inhibited the NGF induction of ODC, as reported by Koizumi et al. (1988), but, surprisingly, potentiated the induction at lower concentrations (2 nM and 20 nM). Staurosporine, at 2 nM, also potentiated the induction by NGF, but this effect was lost at higher concentrations (20 and 200 nM). HBCS, in contrast, potentiated the effects of NGF at all concentrations tested. This striking effect is shown relative to the modest ODC-inducing effects of HBCS alone in Fig. 3.

#### Example 6

The effect of K-252a on choline acetyltransferase (ChAT) activity was assayed in dissociated spinal cord cultures prepared from fetal rats by standard methods (see below). ChAT is the enzyme that catalyzes the synthesis of the neurotransmitter acetylcholine, and is a specific biochemical marker for cholinergic neurons. In the spinal cord, the large majority of cholinergic neurons are motor neurons. Assay of this enzyme may thus be used as an indication of the effects of a factor (or factors) on the survival of cholinergic neurons and/or regulation of this enzyme.

K-252a was added at the indicated concentrations to the cultures after incubating 2-3 hours after plating to allow cells to attach to the substrate.

- 30 -

ChAT activity was measured after 48 hours in culture. K-252a in spinal cord cultures resulted in a dose dependent increase in ChAT activity with maximum efficacy (2- to 3-fold increase) achieved at 200-300 nM (Fig. 4). Higher concentrations resulted in a decrease in ChAT activity (Fig. 4). Longer culture incubation times, up to seven days, resulted in 4-to-5 fold increases in ChAT activity (Fig. 5) due to the decreased basal level of ChAT activity. In this culture system, increasing numbers of motor neurons degenerate and die over time under basal (control) conditions (McManaman et al., Developmental Biol. 125: 311-320, 1988). The results shown in both Figures 4 and 5 are the result of a single application of K-252a on the day of culture initiation, indicating a prolonged effect on the survival of spinal cord cholinergic neurons and/or regulation of the enzyme itself. Methods: Experiments with dissociated cultures of fetal rat spinal cord cells were performed generally as described (Smith et al., J. Cell Biol. 101: 1608-1621, 1985). Dissociated cells were prepared from spinal cords dissected from day 14 embryonic rats by standard techniques known to those skilled in the art, using trypsin dissociation of tissue (Smith et al., 1985). Cells were seeded (plated) at  $6 \times 10^5$  cells/cm<sup>2</sup> in poly-L-ornithine coated plastic tissue culture wells in serum-free N2 medium and incubated at 37°C in a humidified atmosphere of 5% CO<sub>2</sub>/95% air (Bottenstein and Sato, PNAS USA 76: 514-517, 1979) for 48 hours. ChAT activity was measured using modifications of the Fonnum procedure (J. Neurochem. 24: 407-409, 1975) according to Ishida and Deguchi (J. Neurosci. 3: 1818-1823, 1983), and McManaman et al., supra (1988). Activity was normalized to total protein measured by the bicinchoninic acid/Cu<sup>++</sup> reaction (BCA protein assay reagent, Pierce, Rockland, Il).



- 31 -

Example 7

Over one hundred functional derivatives of K-252a were tested in the spinal cord ChAT assay to determine their relative efficacy. The data shown in Figure 8 show that out of the original functional derivatives tested at 300 and 30 nM, 28 resulted in significantly increased ChAT activity at 300 nM. One functional derivative, compound II-21, was also active at 30 nM (30% enhancement of ChAT activity over basal levels). This compound was more potent than K-252a or the remaining analogs since none of these actively enhanced ChAT activity at 30 nM.

Figure 13 shows the effect of the original 28 K-252a derivatives shown to significantly increase ChAT activity in rat spinal cord cultures, as well as 30 additional derivatives (compounds II-29 through II-34, II-36 through II-56, and IV-1 through IV-3, all inclusive).

Example 8

K-252a as well as 50 functional derivatives were assessed for their ability to promote dorsal root ganglion neuron cell survival. Cell survival was measured by uptake of calcein AM, an analog of the viable dye, fluorescein diacetate. Calcein is taken up by viable cells and cleaved intracellularly to fluorescent salts which are retained by intact membranes of viable cells. Microscopic counts of viable neurons correlate directly with relative fluorescence values obtained with the fluorimetric viability assay. This method thus provides a reliable and quantitative measurement of cell survival in the total cell population of a given culture.

Dorsal root ganglion neuron survival was enhanced by K-252a in a concentration-dependent manner; maximum activity was observed at approximately 100 nM (Figure 6). Twenty-four of the 50 analogs tested were

- 32 -

active in promoting DRG neuron survival, twenty-two of which are shown in Figure 7. All of these analogs were also active in increasing spinal cord ChAT activity (see Example 5, Figure 8). The original 22 as well as the 2 additional active analogs (II-30, II-32) are shown in Figure 14. Microscopic examination of the dorsal root ganglion neurons stimulated with the twenty-four active functional derivatives indicated enhanced nerve fiber outgrowth as well.

10           Methods: Dorsal root ganglia were dissected from embryonic age day 8 chick embryos and dissociated cells prepared by subsequent Dispase (neutral protease, Collaborative Research) dissociation. Neurons were seeded at low density ( $1.8 \times 10^4$  cells/cm<sup>2</sup>) into 96 well  
15 poly-L-ornithine and laminin coated plates. Cells were cultured for 48 hours in serum-free N2 medium (Bottenstein and Sato, 1979) at 37°C in a humidified atmosphere, 5% CO<sub>2</sub>/95% air. Cell survival was assessed at 48 hours using the viable fluorimetric assay described  
20 above.

#### Example 9

Infusion of the excitatory amino acid kainic acid (kainate) directly into the ventricles of a rodent brain results in neuronal degeneration of the pyramidal  
25 cells of the hippocampus. This neuronal death is characterized by a marked increase in the proteolysis of the cytoskeletal protein, spectrin. Spectrin breakdown products can be measured in homogenates of the hippocampus within 24 hours following kainate  
30 administration. The magnitude of spectrin proteolysis is highly correlated with the magnitude of neuronal death in pyramidal cells of the hippocampus (Siman et al., J. Neurosci. 9: 1579-1590, 1989), and thus spectrin proteolysis is an excellent biochemical marker of  
35 excitatory amino acid-induced neuronal degeneration.

- 33 -

Excessive release of endogenous excitatory amino acids has been implicated as an etiology in numerous neurological diseases and disorders, including stroke and other ischemic injuries; Alzheimer's disease; motor  
5 neuron disease including amyotrophic lateral sclerosis; Parkinson's disease; Huntington's disease; AIDS dementia; epilepsy; and concussive or penetrating injuries of the brain or spinal cord.

Figure 9 shows the effect of K-252a on  
10 kainate-induced neuronal degeneration in the hippocampus. Cannulated male and female Sprague-Dawley rats received 0.4  $\mu$ g of K-252a, or vehicle, 30 minutes prior to and about 3 and 24 hours following kainate (0.6  $\mu$ g) injection directly into the lateral cerebral ventricles of the  
15 brain (icv). Two weeks later the brains were excised, frozen, sectioned and stained for histological analysis, as described below. Data shown are the mean number of sub-regions of the hippocampus damaged for each group,  $\pm$  S.E.M. K-252a significantly reduced the number of  
20 damaged areas within the hippocampus from  $3.86 \pm 0.78$  (in the absence of K-252a) to  $1.18 \pm 0.4$  (in the presence of K-252a).

Figure 10 shows the effect of K-252a on kainate-induced spectrin breakdown in the hippocampus.  
25 Female Sprague-Dawley rats received 0.4  $\mu$ g of K-252a, or vehicle, together with a neurotoxic dose of kainate (0.6  $\mu$ g), by icv infusion. Sham control animals received infusions of vehicle, but no kainate or K-252a. Twenty-four hours later, homogenates of the dorsal hippocampus  
30 were analyzed for spectrin breakdown products as described below. The magnitude of spectrin proteolysis is expressed as a percent increase in spectrin breakdown products for each group over sham control values. Data shown are the mean percent increase in spectrin breakdown  
35 products for each group (sham = 100%)  $\pm$  S.E.M. Icv

- 34 -

infusion of K-252a significantly reduced the extent of spectrin proteolysis, from about  $140 \pm 15\%$  (in the absence of K-252a) to approximately  $102 \pm 10\%$  (in the presence of K-252a) of sham values.

5

Example 10

Figure 11 shows the effect of HBCS on kainate-induced neuronal degeneration in the hippocampus.

Cannulated female Sprague-Dawley rats received 0.8  $\mu\text{g}$  of HBCS, or vehicle, 40 minutes prior to and about 4 hours  
10 following kainate (0.6  $\mu\text{g}$ ) by icv infusion. Two weeks later the brains were excised, frozen, sectioned and stained for histological analysis, as described below. Data shown are the mean number of sub-regions of the hippocampus damaged for each group,  $\pm$  S.E.M. HBCS  
15 significantly reduced the number of damaged areas within the hippocampus from about  $2.5 \pm 0.6$  (without HBCS treatment) to  $1.3 \pm 0.5$  (with HBCS treatment).

Example 11

Figure 12 compares the effect of three K-252a  
20 functional derivatives on kainate-induced spectrin breakdown in the hippocampus. Female Sprague-Dawley rats received 0.4  $\mu\text{g}$  of K-252a, or compounds III-1, or II-21, or vehicle, together with a neurotoxic dose of kainate (0.6  $\mu\text{g}$ ), by icv infusion. Sham control animals received  
25 infusions of vehicle, but no kainate or K-252a derivative. Twenty-four hours later, homogenates of the dorsal hippocampus were analyzed for spectrin breakdown products as described below. The magnitude of spectrin proteolysis is expressed as a percent increase in  
30 spectrin breakdown products for each group over sham control values. Data shown are the mean percent increase in spectrin breakdown products for each group (sham = 100%)  $\pm$  S.E.M. Icv infusion of K-252a reduced the extent of spectrin proteolysis, from about  $128 \pm 9\%$  (vehicle  
35 treatment) to approximately  $104 \pm 4\%$  (in the presence of

- 35 -

K-252a) of sham values. The K-252a derivatives, III-1 and II-21, failed to prevent kainate-induced spectrin proteolysis.

#### Methods of Figures 9-12

##### 5 Kainate infusion regime:

The effect of K-252a or its derivatives on kainate-induced neuronal damage was evaluated as follows: Adult male or female Sprague-Dawley rats (175-250 g) were anesthetized with Nembutal (50 mg/kg, ip) and  
10 administered a drug, or vehicle treatment, in a total of 5  $\mu$ l, before and after kainate treatment (5  $\mu$ l) by icv infusion using a dose and infusion schedule as indicated for individual cases above. Control animals received vehicle instead of kainate and drug infusion. For  
15 anatomical studies, icv infusions were delivered through a cannula (Plastic One, Roanoke, VA) implanted approximately one week before drug infusions, and positioned at stereotaxic coordinates: anterior-posterior at bregma, 1.5 mm lateral to bregma, and 4.4 mm  
20 ventral from the top of the skull. Results of this treatment regimen were evaluated two weeks later using the anatomical analysis described below.

In studies to assess the effect of K-252a or its derivatives on kainate-induced spectrin proteolysis,  
25 anesthetized rats received a 5  $\mu$ l icv infusion of the drug, or vehicle, simultaneously with kainate, through a 10  $\mu$ l Hamilton syringe positioned at the stereotaxic coordinates described above. These rats were killed 24 hours later and subjected to biochemical analysis as  
30 described below.

#### Anatomical and Biochemical Analyses

Anatomical analysis was performed as follows. Rats were killed by decapitation 2 weeks following treatments, and the brains were rapidly removed and  
35 frozen on dry ice. A series of slide-mounted coronal

- 36 -

sections from each brain was stained with thionin and examined microscopically. Damage to the hippocampus was quantified by summing the total number of 4 anatomically defined regions of the hippocampus (CA1-4 according to the classification of Lorente de No, as described by Shepard, 1979, The Synaptic Organization of the Brain, Oxford, p. 310, hereby incorporated by reference), on both left and right sides of the brain, that suffered a loss of pyramidal cells.

10 Biochemical analysis was performed as follows. Calpain I-sensitive proteolysis of brain spectrin (fodrin) was evaluated in homogenates of the hippocampus using an immunoblot analysis described by Siman et al. (1988, *Neuron*, 1: 279-287, hereby incorporated by reference). Briefly, rats were killed by decapitation 24 hours following treatment, and the dorsal hippocampus was rapidly dissected out of the brain and homogenized in 20 mM Tris-HCl (pH 7.4) containing 0.1 mM phenylmethylsulfonyl fluoride. Proteins from aliquots of each homogenate were separated by SDS-PAGE, and an immunoblot analysis was used to quantitate the amount of kainate-induced spectrin breakdown in each sample.

#### Example 12

K-252a was assessed for its ability to promote survival in striatal cultures. Striata were dissected from embryonic day 17 rat embryos and cells were dissociated by Dispase (neutral protease, Collaborative Research). Neurons were seeded at  $5 \times 10^4$ /well ( $1.5 \times 10^5/\text{cm}^2$ ) in 96-well plates onto wells previously coated with poly-L-ornithine and laminin. Cells were cultured in serum-free N2 medium containing 0.05% bovine serum albumin (Bottenstein and Sato, 1979) at 37°C in a humidified atmosphere, 5% CO<sub>2</sub>/95% air. Cell survival was assessed 5 days after seeding using the calcein viable fluorimetric assay described in Example 8.

- 37 -

Striatal neuronal survival was enhanced by K-252a in a concentration-dependent manner. Maximum activity was found with 75 nM K-252a, and produced an efficacy of 3-4 fold over control (Figure 15). In the control cultures, 90% of the neurons plated on day 0 died within 5 days, whereas in cultures treated with K-252a, 50% of the neurons survived (Figure 16). The survival effect in striatal neurons occurred after 3 days in culture and was sustained for at least 7 days in culture. These results are from a single application of K-252a on the day of culture initiation indicating a sustained effect of survival on a certain population of neurons.

Figure 17 is a pair of photomicrographs taken from control cultures or cultures treated with 75 nM K-252a. There was an increase in cell survival and neurite outgrowth in these cultures in the presence of 75 nM K-252a.

#### Example 13

Thirty-one functional derivatives of K-252a were tested to determine their potency and efficacy in the striatal cell survival assay of Example 10. Figure 18 shows data on 18 K-252a derivatives that promoted the survival of striatal neurons.


#### Processes for Producing Compounds (V)

The processes for producing Compounds (V) are described below.

##### Process 1

Compound (V-1) [Compound V] in which  $R^1$  is  $CH_2SO_2R^7$  and X is  $CO_2R^5$ ] can be prepared by the following reaction step:

- 38 -



(R<sup>5</sup> represents lower alkyl or CH<sub>2</sub>NHCO<sub>2</sub>R<sup>6</sup> in which R<sup>6</sup> represents lower alkyl or aryl; R<sup>7</sup> represents lower alkyl.)

The starting compound (A) is disclosed in Japanese Published Unexamined Patent Application No. 295588/88 (hereby incorporated by reference).


Compound (V-1) can be obtained by treatment of Compound (A) with 1 to 1.5 equivalents of an oxidant. An example of the oxidant is m-chloroperbenzoic acid. As a reaction solvent, a halogenated hydrocarbon such as methylene chloride, chloroform, or ethylene dichloride, or the like is used. The reaction is completed in 0.5 to 1 hour at -20 to 30°C.

### Process 2

Compounds (V-2) [Compound (V) in which R<sup>1</sup> is hydrogen and X is CH<sub>2</sub>NHCO<sub>2</sub>R<sup>6</sup>] can be prepared by the following reaction step:



- 39 -



$R^6$  represents lower alkyl or aryl.

The starting compound (B) is disclosed in Japanese Published Unexamined Patent Application No. 155285/87 (hereby incorporated by reference).

5           Compound (V-2) can be obtained by reaction of Compound (B) with 1 to 3 equivalents  $ClCO_2R^6$  in the presence of 1 to 3 equivalents of a base. An example of the base is triethylamine. As a reaction solvent, a halogenated hydrocarbon such as methylene chloride, 10 chloroform, or ethylene dichloride, or the like is used. The reaction is completed in 0.5 to 3 hours at  $-10$  to  $30^\circ C$ .

#### Example 14

##### Compound II-49

15           Compound (A-1;  $R^5=CH_3$  and  $R^7=C_2H_5$ ) (27 mg, 0.05 mmol) was dissolved in 1 ml of chloroform, and then 10 mg (0.06 mmol) of m-chloroperbenzoic acid was added thereto under ice cooling, followed by stirring at the same temperature for 45 minutes. After dilution with 20 chloroform, the mixture was washed successively with a 8% aqueous solution of sodium thiosulfate, a saturated aqueous solution of sodium bicarbonate, water, and a saline solution, and dried over sodium sulfate. After evaporation of the solvent, the residue was subjected to

SUBSTITUTE SHEET

- 40 -

silica gel column chromatography (chloroform/methanol = 95/5) to give 17.7 mg (yield 62%) of Compound II-49.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ (ppm): 1.298(3H, t, J=7.5Hz),  
2.037 (1H, dd, J=5.0, 14.1Hz),  
5 2.153(3H, s), 3.096(2H, q,  
J=7.5Hz), 3.266 (2H, s), 3.929(3H,  
s), 4.985 (1H, d, J=17.0Hz),  
5.043(1H, d, J=17.0Hz), 6.348(1H,  
s), 7.147 (1H, dd, J=4.9, 7.1Hz),  
10 7.345-8.070(6H, m), 8.612(1H, s),  
9.232(1H, d, J=1.5Hz)

FAB-MS (m/z): 574 (M+1)<sup>+</sup>

#### Example 15

##### Compound II-57

15 Compound (B) (43.8 mg, 0.1 mmol) was dissolved  
in 1 ml of tetrahydrofuran, and then 9.3 μl (0.12 mmol)  
methyl chloroformate and 28 μl (0.2 mmol) of triethylamine  
were added thereto, followed by stirring for 50 minutes  
under ice cooling. After dilution with tetrahydrofuran,  
20 the mixture was washed with a saline solution, and dried  
over sodium sulfate. After evaporation of the solvent,  
the residue was subjected to silica gel column  
chromatography (chloroform/methanol = 99/1) to give 32.6  
mg of Compound II-57.

25 <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ (ppm): 2.099(3H, s), 2.679(1H,  
m), 3.204(1H, dd, J=6.7m 13.8Hz),  
3.837(3H, s), 4.446 (1H, d,  
J=17.3Hz), 4.634 (1H, d,  
J=17.6Hz), 5.497 (1H, brs),  
30 6.591(1H, brs), 7.010-8.037(7H,  
m), 8.592(1H, d, J=6.6Hz)

FAB-MS (m/z): 497 (M+1)<sup>+</sup>

- 41 -

Example 16Compound II-38

Substantially the same procedure as in Example 15 was repeated using 43.8 mg (0.1 mmol) of Compound (B) and 15  $\mu$ l of phenyl chloroformate to give 27.8 mg (yield 50%) of Compound II-38.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm): 2.111(3H, s), 2.890(1H, brd,  $J=13.7\text{Hz}$ ), 3.262(1H, dd,  $J=7.5, 13.9\text{Hz}$ ), 3.742(1H, d,  $J=13.4\text{Hz}$ ), 3.967(1H, d,  $J=12.9\text{Hz}$ ), 4.582(1H, d,  $J=16.3\text{Hz}$ ), 5.342(1H, brs), 5.906(1H, brs), 6.550 (1H, brs), 7.005-8.042(12H, m), 8.596(1H, d,  $J=7.6\text{Hz}$ )

15 FAB-MS (m/z): 559 ( $M+1$ )<sup>+</sup>

Example 17

(The synthesis of Compound H from Compound C is shown in Fig. 19.)

Compound II-39

20 Compound (C) (Japanese Published Unexamined Patent Application No. 295588/88; hereby incorporated by reference) (20 mg, 0.035 mmol) was dissolved in 1 ml of chloroform, and then 14.6  $\mu$ l (0.105 mmol) of triethylamine and 13.9  $\mu$ l (0.175 mmol) of ethyl

25 isocyanate were added thereto, followed by stirring at room temperature for 2 hours. To the solution was added 1 ml of methanol, followed by dilution with chloroform. The mixture was washed successively with water and a saline solution, and dried over sodium sulfate. After

30 evaporation of the solvent, the residue was subjected to silica gel column chromatography (chloroform/methanol = 98/2) to give 21 mg (yield 84% of Compound (D)).

- 42 -

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ (ppm): 1.195(3H, t, J=7.2Hz),  
1.222(3H, t, J=7.2Hz), 1.664(3H,  
s), 2.194(3H, s), 2.555(3H, s),  
3.346(4H, q, J=7.2Hz), 3.820(1H,  
dd, J=7.5, 14.6Hz), 3.938(3H, s),  
5.036(1H, d, J=17.7Hz), 5.125(1H,  
d, J=17.2Hz), 6.745(1H, dd, J=4.8,  
7.4Hz), 7.260-7.898(5H, m),  
8.690(1H, d, J=1.9Hz)

FAB-MS (m/z): 724 (M+1)<sup>+</sup>

Compound (D) (9 mg, 0.012 mmol) was dissolved in a mixture of 0.2 ml of tetrahydrofuran and 0.2 ml of methanol, and then 2 μl of 28% sodium methoxide/methanol was added thereto, followed by stirring at room temperature for 10 minutes. To the solution was added 0.1 ml of a 5% aqueous solution of citric acid, followed by dilution with chloroform. The mixture was washed successively with water and a saline solution, and dried over sodium sulfate. After evaporation of the solvent, the residue was subjected to silica gel column chromatography (chloroform/methanol = 9/1) to give 8 mg of Compound II-39.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ (ppm): 1.086(3H, t, J=7.1Hz),  
1.099 (3H, t, J=7.1Hz), 1.948(1H,  
dd, J=4.8, 14.1Hz), 2.107(3H, s),  
3.158(4H, m), 3.910(3H, s),  
4.880(1H, d, J=17.7Hz), 4.931(1H,  
d, J=16.9Hz), 7.028(1H, dd, J=5.0,  
7.1Hz), 7.332-8.287(5H, m),  
8.838(1H, d, J=2.1Hz)

FAB-MS (m/z): 640 (M+1)<sup>+</sup>

- 43 -

Example 18Compounds II-51 and II-56

Compound (E) (Japanese Published Unexamined Patent Application No. 295588/88; *supra*) (60.7 mg, 0.1 mmol) was dissolved in a mixture of 5 ml of chloroform and 1 ml of methanol, and then 11 mg (0.3 mmol) of sodium borohydride was added thereto under ice cooling, followed by stirring at the same temperature for 15 minutes. After dilution with chloroform, the mixture was washed successively with water and a saline solution, and dried over potassium carbonate. After evaporation of the solvent, the residue was subjected to silica gel column chromatography (Chloroform/methanol/triethylamine = 98/2/0.5) to give 36 mg (yield 59%) of Compound (F).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ (ppm): 1.650(3H, s), 2.027(1H, dd, J=4.9, 14.5Hz), 2.126(3H, s), 3.843(1H, dd, J=7.4, 14.5Hz), 3.891(3H, s), 4.607(2H, s), 4.673(2H, s), 5.125(2H, s), 7.099(1H, dd, J=5.0, 7.3Hz), 7.437-7.907(5H, m), 8.812(1H, d, J=0.8Hz)

FAB-MS (m/z): 612 (M+1)<sup>+</sup>

Compound (F) (159 mg, 0.26 mmol) was dissolved in 15 ml of chloroform, and then 0.8 ml (10.4 mmol) of ethanethiol and 24 mg (0.104 mmol) of camphorsulfonic acid were added thereto, followed by stirring at room temperature for 12 hours. The solution was washed successively with a saturated aqueous solution of sodium bicarbonate, water, and a saline solution, and dried over sodium sulfate. After evaporation of the solvent, the residue was subjected to silica gel column chromatography (ethyl acetate/toluene = 1/9 - chloroform/methanol =

- 44 -

99/1) to give 43 mg of Compound (G) and 75 mg of Compound (H).

Compound (G)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ (ppm): 1.292(3H, t, J=7.4Hz),  
5 1.297 (3H, t, J=7.4Hz), 1.799(3H,  
s), 2.141(1H, dd, J=5.0, 14.5Hz),  
2.256(3H, s), 2.532(2H, q,  
J=7.4Hz), 2.553(2H, q, J=7.4Hz),  
2.869(3H, s), 3.971(1H, dd, J=7.5,  
10 14.5Hz), 3.992(2H, s), 4.005 (3H,  
s), 4.021(2H, s), 5.416(1H, dd,  
J=17.5Hz), 5.459(1H, d, J=17.4Hz),  
6.989(1H, dd, J=5.1, 7.4Hz),  
7.509-7.963(5H, m), 9.134(1H, d,  
15 J=1.2Hz)

FAB-MS (m/z): 700 (M+1)<sup>+</sup>

Compound (H)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ (ppm): 1.294(3H, t, J=7.4Hz),  
1.799(3H, s), 2.149(1H, dd, J=5.0,  
20 14.6Hz), 2.273(3H, s), 2.533(2H,  
q, J=7.4Hz), 2.813 (3H, s),  
3.972(1H, dd, J=7.4, 14.6Hz),  
4.008(3H, s), 4.015(2H, s), 4.951  
(2H, s), 5.377(1H, d, J=17.4Hz),  
25 5.418(1H, d, J=17.4Hz), 6.973(1H,  
dd, J=5.0, 7.5Hz), 7.481-8.037  
(5H, m), 9.093(1H, d, J=1.2Hz)

FAB-MS (m/z): 656 (M+1)<sup>+</sup>

Substantially the same procedure as in Example  
30 17 was repeated using 34 mg of Compound (G) to give 18.7  
mg of Compound II-51.

- 45 -

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ (ppm): 1.300(3H, t, J=7.4Hz),  
1.325(3H, t, J=7.4Hz), 2.185(3H,  
s), 2.514(1H, dd, J=4.8, 14.5Hz),  
2.540(2H, q, J=7.4Hz), 2.555(2H,  
5 q, J=7.4Hz), 3.384(1H, dd, J=7.5,  
14.5Hz), 3.941(2H, s), 3.976(2H,  
s), 4.094(3H, s), 4.836(1H, d,  
J=16.4Hz), 4.910(1H, d, J=16.3Hz),  
5.781 (1H, s), 6.845 (1H, dd,  
10 J=4.8, 7.5Hz), 7.371-7.843(5H,  
m), 8.998(1H, s)  
FAB-MS (m/z): 616 (M+1)<sup>+</sup>

Substantially the same procedure as in Example  
17 was repeated using 30 mg of Compound (H) to give 20.4  
15 mg of Compound II-56.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ (ppm): 1.280(3H, t, J=7.4Hz),  
2.144(3H, s), 2.391(1H, dd, J=4.9,  
14.5Hz), 2.517(2H, q, J=7.4Hz),  
3.320(1H, dd, J=7.4, 14.5Hz),  
20 3.885(2H, s), 4.069(3H, s),  
4.521(1H, d, J=16.3Hz), 4.631(1H,  
d, J=16.7Hz), 4.804(2H, s),  
5.769(1H, s), 6.830(1H, dd, J=4.8,  
7.4Hz), 7.375-7.771(5H, m),  
25 8.934(1H, s)  
FAB-MS (m/z): 572 (M+1)<sup>+</sup>

### Example 19

#### Compound IV-2

Compound (J) (Japanese Published Unexamined  
30 Patent Application No. 120388/87; hereby incorporated by  
reference) (50 mg, 0.09 mmol) was dissolved in a mixture  
of 0.5 ml of trifluoroacetic acid and 50 μl of 3N HCl,

- 46 -

and the solution was stirred at room temperature for 2 days. The precipitates were collected by filtration and subjected to high performance liquid chromatography (Unisil  $\text{C}_{18}$ ; methanol/water = 8/2) to give 8.4 mg of  
5 Compound (IV-2).

$^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ )  $\delta$  (ppm): 4.947 (2H, s),  
7.300-8.010 (6H, m), 8.249 (1H, s),  
9.266 (1H, d,  $J=2.0$  Hz)

FAB-MS ( $m/z$ ): 390 ( $M+1$ )<sup>+</sup>

10

Example 20

Compound II-45 can be prepared by the reaction steps shown in Fig. 20. The starting Compound (J) is disclosed in Japanese Published Unexamined Patent Application No. 120388/87 (hereby incorporated by  
15 reference).

Compound II-45

Compound (J) (200 mg) was dissolved in 1 ml of dimethylformamide, and then 0.25 ml of an aqueous solution of 23.5 mg of sodium hydroxide was added  
20 thereto, followed by stirring at room temperature of 4 hours. After 1N hydrochloric acid was added to adjust the pH of the solution to 1-2, the precipitates were collected by filtration to give 178 mg (yield 91%) of Compound (K).

25

$^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ )  $\delta$  (ppm): 1.965 (1H, dd,  $J=4.8$ ,  
14.0Hz), 2.184 (3H, s), 3.364 (1H,  
dd,  $J=7.5$ , 14.0Hz), 5.029 (1H, d,  
 $J=18.1$ Hz), 5.071 (1H, d,  $J=18.0$ Hz),  
7.133 (1H, dd,  $J=4.9$ , 7.5Hz),  
30 7.595-8.189 (5H, m), 8.733 (1H, s),  
9.398 (1H, d,  $J=2.1$ Hz)

30



- 47 -

Compound (K) (168 mg), was dissolved in 3 ml of pyridine, and then 0.44 ml (4.7 mmol) of acetic anhydride was added thereto, followed by stirring at room temperature for 4 days. After evaporation of the solvent, 4 ml of 1N hydrochloric acid was added to the residue, and the precipitates were collected by filtration to give 182 mg (yield quantitative) of Compound (L).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ (ppm): 1.684(3H, s),  
2.135(1H, dd, J=4.9, 14.4Hz),  
2.252(3H, s), 3.865(1H, dd, J=7.6,  
14.5Hz), 5.063(2H, s), 7.255(1H,  
dd, J=4.9, 7.5Hz), 7.612-8.582(5H,  
m), 8.760(1H, s), 9.389(1H, d,  
J=2.1Hz)

Compound (L) (172 mg) was suspended in thionyl chloride, followed by stirring at 90°C for 4.5 hours. After evaporation of the solvent, diethyl ether was added to the residue and the precipitates were collected by filtration to give 180 mg of Compound (M).

Compound (M) (67 mg, 0.1 mmol) was dissolved in 2 ml of ethylene dichloride, and then 180 μl of aniline in tetrahydrofuran was added thereto under ice cooling, followed by stirring at the same temperature for 1 hour. After evaporation of the solvent, the residue was dissolved in a mixture of 2 ml of tetrahydrofuran and 0.5 ml of methanol, and then 1 ml of 1N NaOH was added thereto, followed by stirring at room temperature for 3 hours. To the solution was added 1N hydrochloric acid (1.2 ml) for neutralization, followed by dilution with tetrahydrofuran. The mixture was washed with a saline solution and dried over sodium sulfate. After evaporation of the solvent, the residue was subjected to

- 48 -

silica gel column chromatography (chloroform/methanol = 98/2) to give Compound II-45 (13 mg from 56 mg of isolated Compound N).

5  $^1\text{H-NMR}$  (DMSO- $d_6$ )  $\delta$  (ppm): 2.110(1H, dd,  $J=4.9$ ,  
13.9Hz), 2.175(3H, s), 5.019(1H,  
d,  $J=18.1\text{Hz}$ ), 5.088(1H, d,  
 $J=18.0\text{Hz}$ ), 6.887(1H, s), 7.119-  
10 8.201(1H, m), 8.711 (1H, s),  
9.391(1H, d,  $J=2.2\text{Hz}$ ), 10.071(1H,  
s)

FAB-MS (m/z): 687 (M+1) $^+$


Other embodiments are within the following  
claims.

What is claimed is:


- 49 -

CLAIMS


1. A composition of the formula



where [Stau] represents a residue of the formula



5 and W represents a radical of the formula



where W' is a hydrocarbylene radical of 2-20 carbon atoms and Y is O or S.

2. A method for enhancing the function of  
10 sensory neurons in a mammal, said method comprising administering to said mammal a therapeutic amount of the composition of claim 1.

3. The method of claim 2, wherein said  
sensory neurons are cholinergic neurons, striatal  
15 neurons, or dorsal root ganglion neurons.

4. A method for treating nerve cell  
degeneration induced by excitatory amino acids, said  
method comprising administering to a mammal a therapeutic  
amount of the composition of claim 1.

20 5. The method of claim 4 wherein said nerve  
cell degeneration is associated with Alzheimer's disease.

- 50 -

6. The method of claim 4 wherein said nerve cell degeneration is associated with motor neuron disease.

7. The method of claim 6 wherein said motor  
5 neuron disease is amyotrophic lateral sclerosis.

8. The method of claim 4 wherein said nerve cell degeneration is associated with Parkinson's disease.

9. The method of claim 4 wherein said nerve cell degeneration is associated with cerebrovascular  
10 disease.

10. The method of claim 9 wherein said cerebrovascular disease is ischemic.

11. The method of claim 4 wherein said nerve cell degeneration is associated with AIDS dementia.

12. The method of claim 4 wherein said nerve  
15 cell degeneration is associated with epilepsy.

13. The method of claim 4 wherein said nerve cell degeneration is associated with concussive injuries to the brain.

14. The method of claim 4 wherein said nerve  
20 cell degeneration is associated with concussive injuries to the spinal cord.


15. The method of claim 4 wherein said nerve cell degeneration is associated with penetrating injuries  
25 to the brain.

- 51 -


16. The method of claim 4 wherein said nerve cell degeneration is associated with penetrating injuries to the spinal cord.

17. The method of claim 4, wherein said nerve cell degeneration is associated with Huntington's disease.


18. A method for enhancing the function of a neuron in a mammal, wherein said neuron is selected from the group consisting of sensory, cholinergic, and striatal neurons, said method comprising administering to said mammal a therapeutic amount of a functional derivative of K-252a, said functional derivative represented by the formula



(II)



(III)



or

15 (IV)

wherein the following substitutions are made:

20	Compound	R <sup>1</sup>	R <sup>2</sup>	X	R	$\begin{matrix} Z^{1(1)} \\ Z^2 \end{matrix}$
	II-1	H	H	CH <sub>2</sub> N <sub>3</sub>	OH	H
	II-2	NHCONHC <sub>6</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-3	CH <sub>2</sub> SOC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
25	II-4	H	H	CH <sub>2</sub> OH	OCH <sub>3</sub>	H
	II-5	H	H	CONHC <sub>2</sub> H <sub>5</sub>	OH	H
	II-6	H	H	CH=NNH-	OH	H
	II-7(2,7)	H	H	CH <sub>2</sub> NH-Gly	OH	H

- 52 -

	II-8	H	H	CON(CH <sub>3</sub> ) <sub>2</sub>	OH	H
	II-9 <sup>(3)</sup>	H	H	-CH <sub>2</sub> NHCO <sub>2</sub> -		H
	II-10	Br	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-11	H	H	CONH <sub>2</sub>	OH	H
5	II-12	H	H	CH <sub>2</sub> OH	OH	H
	III-1	--	H	--	--	H
	II-13	H	H	CONHC <sub>3</sub> H <sub>7</sub>	OH	H
	II-14 <sup>(2)</sup>	H	H	CH <sub>2</sub> NH-Ser	OH	H
	II-15	H	H	CH <sub>2</sub> SOCH <sub>3</sub>	OH	H
10	II-16	H	H	CH=NOH	OH	H
	II-17	H	H	CON O	OH	H
	II-18 <sup>(2,7)</sup>	H	H	CH <sub>2</sub> NH-Pro	OH	H
	II-19	H	H	CH=NNHC(=NH)NH <sub>2</sub>	OH	H
	II-20	Br	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	O
15	II-21	H	H	CONH(CH <sub>2</sub> ) <sub>2</sub> OH	OH	H
	II-22	H	H	CO <sub>2</sub> CH <sub>3</sub>	OH	O
	III-2	--	H	--	--	O
	II-23	H	H	H	OH	H
	II-24	H	H	CH=NNHCONH <sub>2</sub>	OH	H
20	II-25	H	H	CH <sub>2</sub> OCOCH <sub>3</sub>	OH	H
	II-26 <sup>(3)</sup>	H	H	-CH <sub>2</sub> OC(CH <sub>3</sub> ) <sub>2</sub> O-		H
	II-29	NHCONHC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-30	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-31	Br	H	CH <sub>2</sub> OH	OH	H
25	II-32	Br	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-33	CH <sub>2</sub> SC <sub>6</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-34	Cl	Cl	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-36	H	H	CONHC <sub>6</sub> H <sub>5</sub>	OH	H
	II-37	H	H	CH <sub>2</sub> SO	OH	H
30	II-38	H	H	CH <sub>2</sub> NHCO <sub>2</sub> C <sub>6</sub> H <sub>5</sub>	OH	H
	II-39	NHCONHC <sub>2</sub> H <sub>5</sub>	NHCONHC <sub>2</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-40	N(CH <sub>3</sub> ) <sub>2</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-41	CH <sub>3</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-42	CH <sub>2</sub> OCONHC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
35	II-43	NHCO <sub>2</sub> CH <sub>3</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-44	Br	Br	CH <sub>2</sub> OH	OH	H
	II-45	Br	Br	CONHC <sub>6</sub> H <sub>5</sub>	OH	H
	II-46	Br	Br	CONHCH <sub>2</sub> CH <sub>2</sub> OH	OH	H
	II-47	CH <sub>2</sub> OC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
40	II-48	CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-49	CH <sub>2</sub> SO <sub>2</sub> C <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-50	CH <sub>2</sub> S	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-51	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-52	CH=NNH	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
45	II-53	CH <sub>2</sub> S	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-54	CH <sub>2</sub> S(O)	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-55	CH <sub>2</sub> S(O)	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-56	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> OH	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-57	H	H	CH <sub>2</sub> NHCO <sub>2</sub> CH <sub>3</sub>	OH	H

- 53 -


	II-58	Br	H	CONH <sub>2</sub>	OH	H
	II-59	H	H	CH <sub>2</sub> SC <sub>6</sub> H <sub>5</sub>	OH	H
	II-60	H	H	CH <sub>2</sub> S	OH	H
	II-61	H	H	CH <sub>2</sub> SOC <sub>6</sub> H <sub>5</sub>	OH	H
5	IV-1 <sup>(4,9)</sup>	H	H	--	--	H
	IV-2 <sup>(5)</sup>	Br	H	--	--	H
	IV-3 <sup>(6)</sup>	H	H	--	--	H
	IV-4 <sup>(8,9)</sup>	H	H	--	--	H

- 10 (1) Z<sup>1</sup> and Z<sup>2</sup> are both hydrogen, or both are combined together to represent oxygen, where indicated.  
 (2) NH-amino acid linkage is an amide bond through the carboxyl group of the amino acid.  
 (3) X and R are combined together to form the  
 15 linking group.  
 (4) R<sup>3</sup> is CH<sub>2</sub>CH=CH<sub>2</sub>; R<sup>4</sup> is H.  
 (5) R<sup>3</sup> and R<sup>4</sup> are each H.  
 (6) R<sup>3</sup> and R<sup>4</sup> are each CH<sub>2</sub>CH=CH<sub>2</sub>.  
 (7) Compound is in the form of the  
 20 hydrochloride.  
 (8) R<sup>3</sup> is H and R<sup>4</sup> is CH<sub>2</sub>CH=CH<sub>2</sub>.  
 (9) IV-1 and IV-4 is a 1.5 to 1.0 mixture of the two components.

19. The method of claim 18, wherein said  
 25 neuron is a cholinergic neuron.


20. The method of claim 18, wherein said sensory neuron is a dorsal root ganglion neuron, and said functional derivative is represented by the formula (II) or (III):

- 54 -



(II)

or



(III)

wherein the following substitutions are made:

5	Compound <sup>(1)</sup>	R <sup>1</sup>	X	R	Z <sup>1(2)</sup> Z <sup>2</sup>
	II-1	H	CH <sub>2</sub> N <sub>3</sub>	OH	H
	II-2	NHCONHC <sub>6</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-3	CH <sub>2</sub> SOC <sub>2</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
10	II-4	H	CH <sub>2</sub> OH	OCH <sub>3</sub>	H
	II-5	H	CONHC <sub>2</sub> H <sub>5</sub>	OH	H
	II-6	H	CH=NNH-	OH	H
	II-8	H	CON(CH <sub>3</sub> ) <sub>2</sub>	OH	H
	II-9 <sup>(3)</sup>	H	-CH <sub>2</sub> NHCO <sub>2</sub> -		H
15	II-10	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-11	H	CONH <sub>2</sub>	OH	H
	II-12	H	CH <sub>2</sub> OH	OH	H
	III-1	--	--	--	H
	II-13	H	CONHC <sub>3</sub> H <sub>7</sub>	OH	H
20	II-15	H	CH <sub>2</sub> SOCH <sub>3</sub>	OH	H
	II-17	H	CON O	OH	H
	II-19	H	CH=NNHC(=NH)NH <sub>2</sub>	OH	H
	II-20 <sup>(1)</sup>	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	O
	II-21	H	CONH(CH <sub>2</sub> ) <sub>2</sub> OH	OH	H
25	III-2	--	--	--	O
	II-23	H	H	OH	H
	II-24	H	CH=NNHCONH <sub>2</sub>	OH	H
	II-25	H	CH <sub>2</sub> OCOCH <sub>3</sub>	OH	H
	II-30	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	CO <sub>2</sub> CH <sub>3</sub>	OH	H
30	II-32	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	H

(1) R<sup>2</sup> is hydrogen, except in compound II-20 and II-32 where R<sup>2</sup>=Br.

(2) Z<sup>1</sup> and Z<sup>2</sup> are both hydrogen, or both are combined together to represent oxygen, where indicated.

(3) X and R are combined together to form the linking group.



- 55 -

21. The method of claim 2 wherein said composition is administered in conjunction with a neurotrophic factor.

22. The method of claim 4, wherein said composition is administered in conjunction with a neurotrophic factor.

23. The method of claim 18 wherein said functional derivative is administered in conjunction with a neurotrophic factor.


10 24. The method of claim 20 wherein said functional derivative is administered in conjunction with a neurotrophic factor.

25. The method of claims 21, 22, 23, or 24, wherein said neurotrophic factor is a member of the  
15 neurotrophin family.

26. The method of claim 25 wherein said member of the neurotrophin family is nerve growth factor (NGF).


27. The method of claim 18, wherein said neuron is a cholinergic neuron, and said functional derivative  
20 is represented by the formula (II):

- 56 -




wherein  $R^1$  and  $R^2$  are H, X is  $\text{CO}_2\text{CH}_3$ , R is OH, and  $Z^1$  and  $Z^2$  are each H.


28. The method of claim 18, wherein said neuron is a striatal neuron, and said functional derivative is 5 represented by the formula (II), (III), or (IV):



(II)



(III)



(IV)

wherein the following substitutions are made:

	Compound	$R^1$	$R^2$	X	R	$Z^{2(1)}$ $Z^1$
10	K-252a	H	H	$\text{CO}_2\text{CH}_3$	OH	H
	III-1	--	H	--	--	H
	II-1	H	H	$\text{CH}_2\text{N}_3$	OH	H
	II-35	H	H	$\text{CO}_2n\text{-C}_6\text{H}_{13}$	OH	H
15	II-20	Br	Br	$\text{CO}_2\text{CH}_3$	OH	O
	II-10	Br	H	$\text{CO}_2\text{CH}_3$	OH	H
	II-28	$\text{O-n-C}_3\text{H}_7$	H	$\text{CO}_2\text{CH}_3$	OH	H
	II-5	H	H	$\text{CONHC}_2\text{H}_5$	OH	H
	II-29	$\text{NHCONHC}_2\text{H}_5$	H	$\text{CO}_2\text{CH}_3$	OH	H
20	II-2	$\text{NHCONHC}_6\text{H}_5$	H	$\text{CO}_2\text{CH}_3$	OH	H


- 57 -

	II-3	CH <sub>2</sub> SOC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-30	CH <sub>2</sub> SC <sub>2</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-6	H	H	CH=NNH	OH	H
5	II-31	Br	H	CH <sub>2</sub> OH	OH	H
	II-32	Br	Br	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	IV-1 <sup>(2)</sup>	--	H	--	--	H
	II-33	CH <sub>2</sub> SC <sub>6</sub> H <sub>5</sub>	H	CO <sub>2</sub> CH <sub>3</sub>	OH	H
	II-34	Cl	Cl	CO <sub>2</sub> CH <sub>3</sub>	OH	H

- 10 (1) Z<sup>1</sup> and Z<sup>2</sup> are both hydrogen, or both are combined together to represent oxygen, where indicated.  
 (2) R<sup>3</sup> is CH<sub>2</sub>-CH=CH<sub>2</sub>; R<sup>4</sup> is H.

29. The method of claims 19, 27, or 28, wherein said method is used in the treatment of Huntington's  
 15 disease.

30. A composition of the formula (II-4)




II-4

wherein R<sup>1</sup>, R<sup>2</sup>, Z<sup>1</sup>, and Z<sup>2</sup> are each H, X is CH<sub>2</sub>OH, and R is OCH<sub>3</sub>.

- 58 -


31. A composition of the formula (II-14):



II-14

wherein  $R^1$ ,  $R^2$ ,  $Z^1$ , and  $Z^2$  are each H, X is  $\text{CH}_2\text{-NH-Ser}$ , and R is OH.

5            32. A composition of the formula (II-49):




II-49

wherein  $R^2$ ,  $Z^1$ , and  $Z^2$  are each H, R is OH,  $R^1$  is  $\text{CH}_2\text{SO}_2\text{C}_2\text{H}_5$ , and X is  $\text{CO}_2\text{CH}_3$ .

- 59 -


33. A composition of the formula (II-38):



II-38

wherein  $R^1$ ,  $R^2$ ,  $Z^1$ , and  $Z^2$  are each H, R is OH, and X is  $\text{CH}_2\text{NHCO}_2\text{C}_6\text{H}_5$ .

5 34. A composition of the formula (II-45):




II-45

wherein  $R^1$  and  $R^2$  are each Br, R is OH,  $Z^1$  and  $Z^2$  are each H, and X is  $\text{CONHC}_6\text{H}_5$ .

- 60 -


35. A composition of the formula (II-57):



II-57

wherein  $R^1$ ,  $R^2$ ,  $z^1$ , and  $z^2$  are each H, R is OH, and X is  $CH_2NHCO_2CH_3$ .

5                    36. A composition of the Formula (V):



(V)

wherein:

X represents  $CO_2R^5$  or  $CH_2NHCO_2R^6$ ;  
 $R^1$  represents hydrogen or  $CH_2SO_2R^7$ ;  
 $R^5$  represents lower alkyl);  
 $R^6$  represents lower alkyl or aryl; and  
 $R^7$  represents lower alkyl; with the proviso that  
 when  $X = CO_2R^5$ ,  $R^1$  is not hydrogen.

10

FIGURE 1: Effect of K-252a Derivatives on Basal ODC Activity in PC-12 Cells




FIGURE 2: Effect of K-252a Derivatives on NGF-Induced ODC Activity in PC12 Cells






FIGURE 3: Potentiation of NGF Activity by HBCS




FIGURE 4: Effect of K-252a on Choline Acetyltransferase Specific Activity in Rat Spinal Cord Cultures




FIGURE 5: Time Course of K-252a Effect on ChAT Activity in Spinal Cord Cultures






FIGURE 6: Effect of K-252a on Dorsal Root Ganglion Neuron Survival








FIG. 8

FIGURE 9: Effect of K-252a on Kainate-Induced Damage to the Hippocampus




FIGURE 10: Effect of K-252a on Kainate-Induced Spectrin Proteolysis in the Hippocampus




FIGURE 11: Effect of HBCS on Kainate-Induced Damage to the Hippocampus






FIGURE 12: Effects of K-252a Functional Derivatives on  
Kalnate-Induced Spectrin Proteolysis in the Hippocampus




Figure 13: Effect of K-252a and Analogs on ChAT Activity  
In Rat Spinal Cord Cultures: Experimental Table

Compound	Spinal Cord ChAT Activity	
	300nM	30nM
K-252a	100	-
II-44	146	80
II-34	133	66
II-47	126	-
II-1	122	-
II-31	120	-
II-29	118	65
II-32	118	70
II-38	118	-
II-2	111	NT
II-3	109	NT
II-46	109	84
II-5	104	NT
II-41	103	-
II-42	102	64
II-4	100	-
II-30	94	71
II-6	94	-
II-7	90	NT
II-39	90	-
II-36	86	-
II-8	85	-
II-9	83	64
II-10	81	143
IV-3	77	-
II-11	76	NT
II-21	76	63
III-1	75	-
II-12	75	-
IV-1	74	72
II-13	73	-
II-14	71	-
II-15	69	-
II-18	68	-
II-16	68	-
II-17	68	NT
II-40	66	-
II-19	66	-
II-20	65	65
II-45	65	-

Compound	Spinal Cord ChAT Activity	
	300nM	30nM
II-22	62	-
IV-2	62	NT
III-2	60	NT
II-25	58	-
II-37	56	-
II-23	55	-
II-24	50	-
II-26	39	-
II-33	-	77
II-43	-	125
II-49	-	62
II-50	84	56
II-51	130	53
II-52	103	-
II-53	98	67
II-54	51	-
II-55	64	-
II-56	88	53
II-48	70	-

NT = not tested at that concentration

- = not active at that concentration

13/19

Figure 14: Effect of K-252a Analogs on Neuronal Survival in Chick Dorsal Root Ganglion Cultures: Experimental Table

Compound at 300nM	DRG Survival	
	%K-252a	%NGF
III-1	NT	69
III-2	NT	61
II-21	95*	NT
II-10	143	88
II-19	NT	107
II-15	105	158
II-12	83	125
II-17	77	116
II-5	100	110
II-13	73	110
II-9	92	105
II-4	83	95
II-20	55	84
II-1	NT	81
II-2	NT	77
II-23	83	73
II-8	NT	68
II-3	NT	84
II-24	74	67
II-25	68	62
II-11	90	57
II-6	NT	56
II-32	106	NT
II-30	75	NT

NT = not tested

\* = only active at 30nM

Figure 15




Figure 16






Figure 17

EFFECT OF K-252a ON SURVIVAL OF STRIATAL NEURONS *IN VITRO*



CONTROL



75 nM K-252a

17/19

Figure 18 Effect of K-252a and Analogs on Neuronal Survival  
in Rat Striatal Cultures: Experimental Table

Compound	striatal survival % K-252a		
	25nM	75nM	100nM
K-252a		100	
III-1	69	NT	-
II-1	61	NT	104
II-35	52	81	-
II-20	58	-	NT
II-10	91	-	NT
II-28	-	90	-
II-5	104	NT	-
II-29	64	NT	-
II-2	65	-	NT
II-3	50	104	-
II-30	-	114	NT
II-6	81	NT	-
II-31	92	-	NT
II-32	69	NT	-
IV-3	-	NT	47
IV-1	-	NT	66
II-33	-	82	NT
II-34	118	NT	-

NT = not tested at that concentration

- = not active at that concentration








FIG. 20



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US93/06974**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(5) :C07D 498/22, 519/00; A61K 31/40, 31/55

US CL :540/545; 514/211, 410

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 540/545; 514/211, 410

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<u>X</u> Y	US, A, 4,923,986. (Murakata) 8 May 1990, see Col. 1, lines 5-11; Col. 3-4	<u>31</u> 18-20, 23-29
Y	The J. of Antibiotics, Vol. 32, No. 8, issued August 1986, Nakanishi et al., "K-252b, c and d, Potent Inhibitors of Protein Kinase c from Microbial Origin", pgs. 1066-1071. See Fig. 3, pg. 1069.	18-20, 23-29
Y	Chemical Abstracts, Vol. 107, issued 1987, abstract no. 236750y, JP, A, 62,155,284 (Hirata) 10 July 1987.	18-20, 23-29
<u>X</u> Y	Chemical Abstracts, Vol. 107, issued 1987, abstract no. 236751Z, JP, A, 62,155,285 (Hirata et al.), 10 July 1987.	<u>33, 35-36</u> 18-20, 23-29, 32

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be part of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Z" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

05 OCTOBER 1993

Date of mailing of the international search report

01 NOV 1993

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

P. DATLOW ACH

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet)(July 1992)\*

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/06974

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<u>X</u> Y	JP, A, 63,295,588, (Kyowa Hakko Kogyo KK) 01 December 1988, see abstract.	<u>30, 36</u> 18-20, 23-29, 32
Y	Chemical Abstracts, Vol. 111, issued 1989, abstract no. 194456g, JP, A, 63,295,589, (Hirata et al.), 01 December 1988.	18-20, 23-29
Y	Chemical Abstracts, Vol. 107, issued 1987, Abstract No. 197943s, JP, A, 62, 120, 388, (Akiyama et al.) 01 June 1987.	18-20, 23-29
Y	Annals of the N.Y. Acad. of Sciences, Vol. 559, issued 05 July 1989, Abe et al., "Arachidonic Acid Metabolism in Ischemic Neuronal Damage", pgs. 259-268, see pgs. 260, 264 and 267.	18-20, 23-29
Y	Neuroscience Letters, Vol. 108, issued 01 January 1990, G.D. Borasio, "Differential effect of the protein kinase inhibitor K-252a on the in vitro survival of chick embryonic neurons", pgs. 207-212, see abstract pg. 207.	18-20, 23-29
Y	The J. of Pharmacology and Exp. Therapeutics, Vol. 257, No. 2, issued 02 January 1991, Nabeshima et al., "Staurosporine facilitates recovery from the Basal Forebrain-Lesion-Induced Impairment of Learning and Deficit of Cholinergic Neurons in Rats", pgs. 562-566., see pg. 566, first ten lines.	18-20, 23-29

Form PCT/ISA/210 (continuation of second sheet)(July 1992)\*

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/06974

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

(Telephone Practice)

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet(1))(July 1992)★

**B. FIELDS SEARCHED**

Electronic data bases consulted (Name of data base and where practicable terms used):

STN-Registry File, HCA File, CA previews File, based upon compound structure and utility.

APS-Text-"PKC or Protein Kinase Inhibitor" and "Nerve" or "Neuron".

**BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING**

This ISA found multiple inventions as follows:

Group I, claims 1-17, 21-22, 25-26, drawn to Bis-Stanrosporine derivatives, compositions and methods of treatment.

Group II, claims 18-20, and 23-36, drawn to K-252a derivatives of formula II and method of treatment.

Group III, claims 18-20, 23-26, and 28-29, drawn to method of treatment using K2529 derivatives of formula III or IV.

The inventions listed as Groups I-III do not meet the requirements for Unity of Invention for the following reasons:

The three groups of inventions are based upon structurally different compounds such that a reference which anticipates one group would not make obvious the other groups. Each group would raise different issues of novelty, inventive step and industrial applicability.

